

Mammalian Prion Biology: One Century of Evolving Concepts

Review

Adriano Aguzzi* and Magdalini Polymenidou

Institute of Neuropathology
University Hospital of Zürich
Schmelzbergstr. 12
CH-8091 Zürich
Switzerland

Prions have been responsible for an entire century of tragic episodes. Fifty years ago, kuru decimated the population of Papua New Guinea. Then, iatrogenic transmission of prions caused more than 250 cases of Creutzfeldt-Jakob disease. More recently, transmission of bovine spongiform encephalopathy to humans caused a widespread health scare. On the other hand, the biology of prions represents a fascinating and poorly understood phenomenon, which may account for more than just diseases and may represent a fundamental mechanism of crosstalk between proteins. The two decades since Stanley Prusiner's formulation of the protein-only hypothesis have witnessed spectacular advances, and yet some of the most basic questions in prion science have remained unanswered.

Introduction

A few years ago, it was memorably stated that prion diseases (also termed transmissible spongiform encephalopathies, or TSE) constitute one of the best-understood groups of neurodegenerative diseases (DeArmond and Prusiner, 1995). Depending on whom you ask, this statement may be regarded as entirely correct or completely off the mark.

Of course, prion diseases are quite well understood. Largely thanks to the enthusiasm and intuition of pioneers such as Stanley Prusiner and Charles Weissmann, progress in prion science has experienced two decades of quantum leaps. These include the isolation of the disease-associated, protease-resistant prion protein, PrP^{Sc} (Bolton et al., 1982), the formulation of the protein-only hypothesis (Prusiner, 1982), the cloning of the *Prnp* gene that encodes PrP^C and the startling realization that it is a normal, cellular gene (Basler et al., 1986; Chesebro et al., 1985; Oesch et al., 1985), the discovery that the host-determined aspects of the “species barrier” are crucially governed by the sequence of PrP^C (Scott et al., 1989), the linkage between *PRNP* mutations and hereditary prion disease (Hsiao et al., 1989), and the demonstration that PrP^C-deficient mice are alive and well, but resistant to prion diseases (Büeler et al., 1992, 1993).

And yet one may argue that prions are not well understood at all! We are still unable to precisely pinpoint the physical nature of the agent (Chesebro, 1998), and we do not avail of any high-resolution molecular structure of PrP^{Sc}. Hence, the models of conversion of PrP^C to PrP^{Sc} are speculative at best, and the conversion pro-

cess could not be reproduced under cell-free conditions in a way that would lead to replication of prion infectivity. Finally, precious little knowledge is available on how the infectious agent damages the brain, and the function of the normal protein continues to be obscure.

The Timeline of TSE Research

In one or the other form, prions have captured a sizeable mind share for almost two centuries (Table 1). Scrapie—the prototypic prion disease affecting sheep and goat—had been a concern since the 19th century. This is understandable given the importance of the wool textile business in the industrial revolution. But the crucial breakthrough was already achieved in the 1930s by the experimental transmission of scrapie to goats (Cuille and Chelle, 1939). Little happened in the two following decades, until Carleton Gajdusek showed that kuru, which was decimating the aborigines of Papua New Guinea (Gajdusek and Zigas, 1957), was a transmissible spongiform encephalopathy. Interestingly, the first attempts at transmitting kuru to primates failed for the same reason that experimental transmission of scrapie among sheep had failed for decades: the incubation time of the disease was longer than the patience of the investigators (Schwartz, 2003). Following a concise suggestion by William Hadlow that kuru resembled scrapie, and hence might exhibit a very long incubation time (Hadlow, 1959), Gajdusek achieved transmission of kuru to chimps (Gajdusek et al., 1966, 1967) and, shortly thereafter, transmission of Creutzfeldt-Jakob disease (CJD) (Gibbs et al., 1968).

It is remarkable (and somewhat sobering) to note that some of the questions that had already been formulated in the 19th century are still open. For example, is sheep scrapie a predominantly genetic or infectious disease? If the latter is true, how does it spread among flocks? The wildfire-like epizootic of chronic wasting disease in North American cervids (Williams and Young, 1980), as well as the “scrapie eradication plan” of the European Union (which aims at selective breeding of purportedly scrapie-resistant sheep genotypes), bears the most recent witness to the general importance of these issues.

The Nature of the Prion

Throughout this paper, the term “prion” is used to denote the infectious principle active in TSEs. The various hypotheses of TSE pathogenesis state that the prion may be congruent, partially overlapping, or different from the protease-resistant form of PrP found in prion diseases, which is termed PrP^{Sc}.

Two papers reprinted in the current issue of *Cell* represent two major turning points in prion research. The first paper describes the discovery, by Stanley Prusiner and coworkers, of a crucial property of the prion: its remarkable resilience against proteolytic degradation (McKinley et al., 1983). Digestion with 50 μg/ml of proteinase K (PK) at 37°C for 2 hr would not degrade the carboxy-proximal domain of PrP^{Sc} nor decrease the infectious titer of the prion preparation. But PrP^{Sc} is not “unbreakable” and can eventually be digested by more vigorous enzymatic treatment—in which case prion infectivity titers will also subside. This remarkable discovery identi-

*Correspondence: adriano@pathol.unizh.ch

Table 1. Essential Chronology of Prion Research

Mid 18 th Century	Earliest Description of Scrapie Recorded
1898	Neuronal vacuolation discovered in brains of scrapie-sick sheep
1918	Contagious spread of scrapie in natural conditions suspected
1920	First cases of CJD described (Creutzfeldt, 1920; Jakob, 1921)
1937	Scrapie epidemic in Scotland following administration of formalin-treated louping ill vaccine prepared from sheep brain
1939	Experimental transmission of scrapie reported (Cuille and Chelle, 1939)
1955–1957	Kuru discovered among Fore people of Papua New Guinea (Gajdusek and Zigas, 1957)
1959	Similarities between kuru and scrapie noted (Hadlow, 1959)
1961	Multiple strains of scrapie agent described (Pattison and Millson, 1961)
1961	Scrapie transmitted to mice (Chandler, 1961)
1963	Transmission of kuru to chimpanzees reported (Gajdusek et al., 1966)
1966	Scrapie agent found to be highly resistant to ionizing radiation and ultraviolet light (Alper et al., 1966, 1967)
1967	First enunciation of the protein-only hypothesis (Griffith, 1967)
1968	CJD transmitted to chimpanzees (Gibbs et al., 1968)
	Description of <i>Sinc</i> gene affecting scrapie incubation period in mice (Dickinson et al., 1968)
1974	First documented iatrogenic prion transmission (corneal graft) (Duffy et al., 1974)
1980	Protease resistant, highly hydrophobic protein discovered in hamster brain fractions highly enriched for scrapie infectivity (Prusiner et al., 1980)
1982	Prion concept enunciated (Prusiner, 1982)
1985	Gene encoding PrP ^C cloned (Chesebro et al., 1985; Oesch et al., 1985)
1986	PrP ^C and PrP ^{Sc} isoforms shown to be encoded by same host gene (Basler et al., 1986)
1987	Linkage between <i>Prnp</i> and scrapie incubation period in mice (Westaway et al., 1987)
	First report of BSE in cattle (Wells et al., 1987)
1989	Mutation in PrP linked to Gerstmann-Sträussler syndrome (Hsiao et al., 1989)
	Importance of isologous PrP ^C /PrP ^{Sc} interactions established (Scott et al., 1989)
1992	Ablation of <i>Prnp</i> by gene targeting in mice (Büeler et al., 1992)
1993	<i>Prnp</i> ^{0/0} mice are resistant to scrapie inoculation (Büeler et al., 1993; Sailer et al., 1994)
	Structural differences between PrP ^C and PrP ^{Sc} isoforms noted (Pan et al., 1993)
1994	Cell-free conversion of PrP ^C to protease-resistant PrP (Kocisko et al., 1994)
1996	New variant of CJD identified (Will et al., 1996)
	BSE prion strain carries a distinct glycoctype signature (Collinge et al., 1996)
	First NMR structure of core murine PrP ^C solved (Riek et al., 1996)
1997	Evidence that nvCJD is caused by the BSE agent (Bruce et al., 1997; Hill et al., 1997a)
	B lymphocytes necessary for peripheral prion pathogenesis (Klein et al., 1997)
1998	Genes controlling incubation period are congruent with <i>Prnp</i> (Moore et al., 1998)
1999	Discovery of the PrP ^C homolog (Moore et al., 1999)
2000	Temporary depletion of lymphoid FDCs impairs prion replication (Montrasio et al., 2000)
	Experimental transmission of BSE in sheep by blood transfusion (Houston et al., 2000)
2001	Complement involved in prion pathogenesis (Klein et al., 2001; Mabbott et al., 2001)
2003	Transgenic expression of soluble PrP inhibits prion replication (Meier et al., 2003)

fied PrP^{Sc} as the first reliable surrogate marker of prion infection. The impact of this technology was phenomenal: even now, twenty years after its original description, the detection of PK-resistant prion protein (termed PrP²⁷⁻³⁰ because of its molecular weight after hydrolysis of its PK-sensitive amino-terminal domain) remains the gold standard for biochemical diagnosis of prion diseases and forms the basis for all of the currently marketed BSE tests (Figure 1).

The second paper, to which one of us had the privilege of contributing, verifies a crucial prediction of Prusiner's protein-only hypothesis (Büeler et al., 1993). If PrP^{Sc} multiplies by imparting its conformation onto host-borne PrP^C, organisms devoid of PrP^C should be resistant to prion infection. This idea was compelling, but in the early days of prion research, no technology was available that would allow for the targeted removal of a specific gene from the mammalian genome. As soon as *in vivo* gene ablation became feasible (Zijlstra et al., 1990), Hansrüedi Büeler and Charles Weissmann set out to ablate the *Prnp* gene, which encodes PrP^C. *Prnp*^{0/0} mice were alive and well (Büeler et al., 1992), notwithstanding some minor abnormalities (Collinge et al., 1994; Tobler et al., 1996; Watarai et al., 2003)—some of which may not even be causally related to the prion gene (Aguzzi and Hardt,

2003). The excitement in Zurich was considerable as it became gradually clear that inoculation of *Prnp*^{0/0} mice with brain homogenate from scrapie-sick mice failed to induce disease of any kind (Büeler et al., 1993) or elicit any subclinical replication of the agent (Sailer et al., 1994).

The study of Büeler and colleagues has sometimes been invoked as the “final proof” of the protein-only hypothesis. That is certainly not the case: the knockout experiment was designed to *disprove* Prusiner's hypothesis—and it would have certainly done so if *Prnp*^{0/0} mice had developed disease. As always with negative results, alternative interpretations can be offered (Popper, 1991). Those skeptical of the prion hypothesis were quick in pointing out that PrP^C may be a receptor for a hitherto unidentified virus, whose ablation would confer antiviral resistance. Yet it is fair to say that the resistance to scrapie of *Prnp* knockout mice constitutes one of the most stringent challenges to the protein-only hypothesis. Hence its failure is very significant.

The availability of *Prnp*^{0/0} mice has triggered a cascade of technological and conceptual advances. For example, it emerged that PrP^C, besides controlling prion replication, is necessary for neuronal damage: *Prnp*^{0/0} neurons adjacent to infected *Prnp*^{+/+} brain grafts do not




Figure 1. Diagnostic Procedures for Prion Diseases

Synopsis of current diagnostic methods for TSE in humans (left panel) and in experimental animals (right panel). Most methods rely upon the detection of PK-resistant PrP^{Sc}. The tissues in which PrP^{Sc} has been detected in humans are listed in the middle panel. While PrP^{Sc} deposits are most abundant in the CNS, the list of peripheral organs in which PrP^{Sc} can be detected has significantly grown in recent years; it now includes most lymphoid organs as well as skeletal muscle.

incur damage (Brandner et al., 1996a). PrP^C is also involved in the transport of the infectious agent from peripheral sites to the central nervous system: its expression appears to be needed in a sessile compartment (Blättler et al., 1997), which is likely to be congruent with stromal components of the lymphoreticular tissue (Montrasio et al., 2000) and of the peripheral nervous system (Glatzel et al., 2001). The microenvironment of lymphoid organs appears to control the velocity of neuroinvasion (Prinz et al., 2003a).

PrP^C is not only produced by neurons; its expression is in fact quite ubiquitous, notably including lymphocytes (Cashman et al., 1990) and stromal cells of lymphoid organs (Kitamoto et al., 1991). As a result, wild-type mice enjoy an extremely tight immunological tolerance against PrP^C, which had rendered the production of high-affinity immunoreagents very difficult. Instead, the immunization of Prnp^{0/0} mice yielded large numbers of very high-affinity antibodies, some of which form the basis for the current crop of BSE tests.

Still, it proved difficult to generate conformational anti-

bodies discriminating between PrP^C and PrP^{Sc}. This is surprising in view of the dramatic structural differences between these two isoforms and their differential binding to serum proteins (Fischer et al., 2000). Does the failure of the immune system to generate antibodies specific for PrP^{Sc} indicate that all relevant neopeptides of PrP^{Sc} that are newly exposed by the conversion of the protein to its disease-associated state are inaccessible? Early claims of discriminatory antibodies, such as Prionics' 15B3 clone (Korth et al., 1997), have not lived up to the expectations. A recently developed antibody against a characteristic tripeptide (YYR) exposed in PrP^{Sc}, but not in PrP^C, may be more promising (Paramithiotis et al., 2003). However, the YYR motif is certainly not specific to PrP^{Sc}, and the usefulness of this antibody awaits independent confirmation.

The Spontaneous Generation of Prions

A mesmerizing implication of the protein-only hypothesis is the propagation of prions in an entirely synthetic system. If the infectious agent is a misfolded form of PrP, and its replication is promoted by its interaction

with PrP^C, then the entire process should be, in principle, reproducible in a cell-free environment consisting exclusively of PrP^C, PrP^{Sc}, and maybe some “promoting factors.” The importance of such an experiment is immediately evident: *de novo* generation or amplification of prions from defined components would prove the protein-only hypothesis and set to rest all other alternative explanations (Aguzzi and Weissmann, 1997). Besides, such a system would be extremely valuable for studying the conversion process, for exploring the species barrier phenomenon, and for testing conversion antagonists that may provide therapeutic compounds.

A decisive milestone toward this goal was accomplished with the establishment of an *in vitro* conversion system based on the coincubation of substantially purified constituents (Kocisko et al., 1994). This seminal work showed that incubation of radiolabeled PrP^C with cold PrP^{Sc} leads to the formation of PK-resistant radiolabeled PrP—indicating that PrP^{Sc} had somehow imparted some of its properties onto PrP^C. The original system required vastly superstoichiometric amounts of PrP^{Sc}, which precluded the detection of any increase in prion infectivity. However, the method was used to probe the conversion efficiency between PrP molecules with different primary sequences and thereby, to some extent, the tightness of species barriers (Bessen et al., 1995; Horiuchi et al., 2000). In the intervening years, the *in vitro* conversion methodology has yielded remarkable insights and even assays for identification of anti-prion compounds.

It was reported that PrP^{Sc} could be amplified by cycles of sonication followed by incubation with brain homogenate (Saborio et al., 2001). The idea behind this experiment was that sonication might fracture large PrP^{Sc} aggregates into smaller units, each one of which would accrue PrP^C and act as independent “infectious unit.” Several skeptics, however, have pointed out that this intriguing report is still awaiting independent confirmation. Also, more than two years after its publication, no evidence has come forward that this “protein misfolding cyclic amplification” would augment the infectivity of any given sample.

Along parallel lines, conditions were established at that recombinantly produced PrP was transformed into an isoform termed β PrP, with several typical properties of PrP^{Sc} (Jackson et al., 1999): increased β sheet content, aggregability, and resistance to PK. This molecule was deemed quite interesting for two main reasons. Firstly, one had hoped that immunization of mice with β PrP might give rise to conformation-specific monoclonal antibodies, which would help in discriminating directly between PrP^C and PrP^{Sc}. The latter would render obsolete the venerable PK digestion assay and may facilitate the development of higher-throughput PrP^{Sc} immunoassays. Secondly, β PrP might be equivalent to PrP* (Weissmann, 1991), a metastable intermediate postulated to arise during conversion of PrP^C to PrP^{Sc}. If so, inoculation of mice with suitable amounts of β PrP might result in the generation of transmissible disease.

At present β PrP has yet to fulfill either of these two expectations. Yet it is not implausible that additional experimentation in specifically devised animal models may change this negative outcome. As with transmission of kuru to chimps, it is advisable to be patient.

Form Follows Function

If the protein-only hypothesis is correct, one could argue that the prion problem is, in essence, one of protein structure. Whether prions multiply by template-directed refolding or by seeded nucleation, certain domains of PrP^C (or the entire protein) would need to rearrange such that the monomeric protein becomes capable of inducing the same change in further PrP^C monomers (Figure 2A). This idea represents the core of the “template-directed refolding” hypothesis, which predicates an instructionist role for PrP^{Sc} onto PrP^C. The experimental evidence is compatible with this hypothesis, yet no positive evidence in its favor has come forward.

Alternatively, it has been proposed that PrP^{Sc} exists in a mass-action equilibrium with PrP^C. Such equilibrium would be heavily shifted toward the side of PrP^C so that only minute amounts of PrP^{Sc} would coexist with PrP^C. If that were the case, PrP^{Sc} could not possibly represent the infectious agent since it would be ubiquitous. According to this “nucleation” hypothesis (Jarrett and Lansbury, 1993), however, the infectious agent would consist of a highly ordered aggregate of PrP^{Sc} molecules. The aggregated state would be an intrinsic property of infectivity: monomeric PrP^{Sc} would be harmless, but it might be prone to incorporation into nascent PrP^{Sc} aggregates (Figure 2B).

Testing these hypotheses requires precise knowledge of the structural features of both PrP^C and PrP^{Sc}. To date, such knowledge has not progressed to a state that would allow for resolution of this question. The structure of PrP^C has been studied extensively with high-resolution methods. Both crystallography (Knaus et al., 2001) and nuclear magnetic resonance (NMR) spectroscopy (Riek et al., 1996) have yielded detailed insights into the arrangement of PrP^C at the atomic level. PrP^{Sc}, however, has been amenable merely to low-resolution structural methods.

The NMR studies of recombinant PrP^C yielded a big surprise. The amino-proximal half of the molecule is not structured at all, whereas the carboxy-proximal half is globular and contains three α helices (Riek et al., 1996, 1997). This does not mean that the amino terminus must be randomly coiled *in vivo*: functional studies in transgenic mice imply that the domain comprising amino acids 32-121 carries out important physiological functions (Shmerling et al., 1998). Maybe the flexible tail of PrP^C acquires a defined structure once it reaches its natural habitat on rafts, which are specialized microdomains of the plasma membrane (Naslavsky et al., 1997).

Why wasn't it yet possible to elucidate the structure of PrP^{Sc}? As discussed above, prion infectivity can be recovered only from prion-infected mammalian organisms or (in much lesser quantities) from infected cultured cells. In neither case is the purity of the recovered material satisfactory. Moreover, infectivity-associated PrP^{Sc} appears to consist obligatorily of aggregates; disaggregation sterilizes prions (Prusiner et al., 1981). But insoluble aggregates are resilient to most technologies for determination of protein structure; hence all we know is that PrP^{Sc} consists mainly of β -pleated sheet (Caughey et al., 1991) and that PrP^{Sc} aggregates expose a remarkably ordered structure (Wille et al., 2002).




Figure 2. Models for the Conformational Conversion of PrP^C into PrP^{Sc}

(A) The “refolding” or template assistance model postulates an interaction between exogenously introduced PrP^{Sc} and endogenous PrP^C, which is induced to transform itself into further PrP^{Sc}. A high energy barrier may prevent spontaneous conversion of PrP^C into PrP^{Sc}.

(B) The “seeding” or nucleation-polymerization model proposes that PrP^C and PrP^{Sc} are in a reversible thermodynamic equilibrium. Only if several monomeric PrP^{Sc} molecules are mounted into a highly ordered seed, further monomeric PrP^{Sc} can be recruited and eventually aggregates to amyloid. Within such a crystal-like seed, PrP^{Sc} becomes stabilized. Fragmentation of PrP^{Sc} aggregates increases the number of nuclei, which can recruit further PrP^{Sc} and thus results in apparent replication of the agent.

Yeast Prions

Thirty years ago, Francois Lacroute described mysterious yeast traits that apparently propagated by nonmendelian genetics (Lacroute, 1971). For two decades, this phenomenon remained unexplained—until Reed Wickner proposed that the unusual genetic properties of these mutants could be explained by a prion-like behavior of two previously identified yeast proteins: Sup35p, an essential component of the translation termination machinery, and Ure2p, a protein that regulates nitrogen metabolism (Wickner, 1994). Further yeast prions were identified in the following, so that one could now argue that the yeast prion phenomenon is much better understood than its mammalian counterpart. The prion-forming domain (PrD) of Sup35p is modular and transferable; artificial prions were generated by fusing a mammalian receptor to the Sup35p PrD (Li and Lindquist, 2000).

In the prion-infected state (termed ψ^+), Sup35p is sequestered into fibrils. As consequence, termination of translation is impaired, and reading frames situated downstream of nonsense codons can be translated into proteins (Figure 3). Just like in street traffic, ignoring stop signs does not generally constitute healthy behavior, but Susan Lindquist made a convincing case that such transgressions may play a decisive role in creating “evolutionary buffers.” By occasionally switching on bicistronic reading frames through the ψ^+ state, yeast cells can reversibly probe the effects of combinatorial expression of mutated genes, hence creating additional layers of evolutionary variation (True and Lindquist, 2000).

BSE and Other Prion Threats to Humans

When Stanley Prusiner started his first attempts at tackling the problem of TSE (Prusiner et al., 1977), this group of diseases was not exactly in the public limelight. However, bovine spongiform encephalopathy (BSE) was recognized a few years later (Wells et al., 1987) —an event that would dramatically change the public perception of prion diseases. CJD was, and fortunately continues to be, exceedingly rare: its incidence is typically 1/10⁶ inhabitants/year, but reaches 3/10⁶ inhabitants/year in Switzerland, which is currently reporting the highest number of cases (Glatzel et al., 2002, 2003b). Kuru, once decimating the population of Papua New Guinea, has almost disappeared. Iatrogenic transmission of CJD has principally occurred through improperly sterilized neurosurgical instruments, transplants of dura mater, and administration of pituitary hormones of cadaveric origin. While the two latter routes of transmission no longer pose a major threat, a significant number of individuals may have been infected during a critical time window and may develop CJD in the coming years.

Variant CJD (vCJD) has caused some 140 deaths in the United Kingdom and a few cases in France, Italy, and Canada (<http://www.doh.gov.uk/cjd/stats/aug02.htm>). Epidemiological, biochemical, and histological evidence suggests that vCJD represents transmission of bovine spongiform encephalopathy (BSE) prions to humans (Aguzzi, 1996; Aguzzi and Weissmann, 1996; Bruce et al., 1997; Hill et al., 1997a). The incidence of vCJD in the United Kingdom rose each year from 1996 to 2001, evoking fears of a large upcoming epidemic.




Figure 3. Function of the Yeast Prion, Sup35
(A) Sup35 consists of an amino-terminal glutamine-rich module crucial for conversion into the prion state.
(B) In the Ψ^- state, Sup35 is required for reliable termination of translation.
(C) In Ψ^+ yeast cells, however, Sup35 is sequestered in ordered fibrillary aggregates. Shortage of functional Sup35 leads to transgressions in stop codon recognition and translation of downstream reading frames (red line). In the off state, such pseudogenes may accumulate otherwise toxic mutations. Acquisition of the on (Ψ^+) state may lead to the appearance of new phenotypes, hence increasing the complexity of genetic variability.

Since the year 2001, however, the incidence of vCJD in the UK appears to be stabilizing (<http://www.cjd.ed.ac.uk/vcjdq.htm>). One may argue that it is too early to draw any far-reaching conclusions, but each year passing without any dramatic rise in the number of cases increases the hope that the total number of vCJD victims will be limited (Valleron et al., 2001). Presently, there is reason to hope that the incidence of vCJD in the United Kingdom may already be subsiding (Andrews et al., 2003).

vCJD prions accumulate prominently in lymphoreticular tissue, and the latter can be used for diagnostic purposes. Surprisingly, prions accumulate in lymphoid organs and muscle of sporadic CJD patients (Glatzel et al., 2003a).

There is uncertainty surrounding the danger of transmission to humans represented by chronic wasting disease. In fact, even transmissibility of BSE to humans relies on circumstantial evidence. Epidemiology and biochemistry favor the link between BSE and vCJD, but are not ultimately conclusive. The Koch postulates (which would unambiguously assign an infectious agent to a disease) have never been fulfilled, and experimental inoculation of humans was fortunately never performed. Also, accidental exposure to BSE infectivity of a sizable collective at a precisely defined time point has never occurred, or did not result in disease. Likewise, we do not know whether scrapie is just a veterinarian problem that affects only sheep and goat or whether it can cross species barriers and affect humans. Finally, it is unknown whether BSE, upon transmission to sheep, remains as dangerous for humans as cow-derived BSE, or whether it becomes attenuated and acquires the (allegedly) innocuous properties of bona fide sheep scrapie.

The Elusive Function of PrP^C

In spite of the fact that the first *Prnp* knockout mice are available since 12 years (Büeler et al., 1992), the normal

function of the cellular prion protein is still unknown. A number of subtle abnormalities have been described in PrP-deficient mice (Collinge et al., 1994; Tobler et al., 1996), but their molecular basis is undefined, and there may be some variability due to the genetic background of the mice utilized. Hence, the only definite phenotype of *Prnp*^{0/0} mice is their resistance to prion inoculation (Büeler et al., 1993) —yet it seems unlikely that a singular protein that is as highly conserved among species as PrP^C, from turtles to frogs, fish, and humans, has evolved for the sole reason of bestowing susceptibility to prion diseases.

If the function of PrP^C were completely unrelated to prion disease pathogenesis, one might argue that PrP^C is just one of many thousands proteins whose function awaits clarification —but why should then the elucidation of the function of PrP^C be given any priority? On the other hand, the function of PrP^C may very well have something to do, in a subtle way, with prion-induced damage. *Prnp* ablation does not elicit disease, even when induced postnatally (Mallucci et al., 2002); hence prion pathology is unlikely to come about by loss of PrP^C function. But assume that PrP^C transduces a signal, or that it possesses some enzymatic activity. If so, conversion to PrP^{Sc} may alter signal transduction strength, or substrate specificity, thereby conferring a toxic dominant function. In these scenarios, understanding the function of PrP^C may help in deciphering prion pathology and maybe even devising therapeutical approaches.

So, what is the evidence that PrP^C may be a signal transducer or an enzyme? Speculations on both hypotheses abound, but facts are scarce. Crosslinking PrP^C with F(ab)₂ antibody fragments has been reported to activate intracellular tyrosine kinases (Mouillet-Richard et al., 2000). However, this phenomenon was not reported to occur in vivo, and the only cell line in which it was described was never made available to the scientific community for independent verification. This does not exclude that PrP^C functions as a signal transducer, but the present case is of limited strength.

Is PrP^C an enzyme? Glockshuber noted that PrP^C has similarities to membrane-anchored signal peptidases (Glockshuber et al., 1998), but his observation has not been substantiated by functional data. The speculation that PrP^C may be a superoxide dismutase (Brown et al., 1997, 1999) was perceived as particularly attractive in view of its multiple copper binding sites, and it was recently suggested that amino-proximally truncated PrP^C may depress endogenous dismutase activity (Sakudo et al., 2003). However, PrP^C does not make any measurable contribution to dismutase activity in vivo (Hutter et al., 2003; Waggoner et al., 2000).

Maybe PrP^C and PrP^{Sc} do not possess any intrinsic biological activity, yet they modify the function of other proteins. This supposition has prompted a search for PrP-interacting partners, and there is no dearth of PrP binding proteins: the antiapoptotic protein Bcl-2 (Kurschner et al., 1995), caveolin (Gorodinsky and Harris, 1995; Harmey et al., 1995), the laminin receptor precursor (Rieger et al., 1997), plasminogen (Fischer et al., 2000), and N-CAM (Schmitt-Ulms et al., 2001). None of these interactors, however, have yet revealed a functional pathway in which PrP^C would be involved in vivo. It was recently shown that PrP-deficient macrophages do not support bacterial “swimming internalization” of the Gram-negative bacterium, *Brucella abortus* (Watarai et al., 2003), and that PrP^C interacts with a *Brucella* heat shock protein, Hsp60. These findings raise the question of whether PrP^C may participate in a general Hsp60-dependent “danger sensing” mechanism (Aguzzi and Hardt, 2003).

A Doppelganger of the Prion Protein

The original *Prnp*^{0/0} mice did not display any severe abnormalities. However, some of the knockout lines generated later, i.e., Nsgk *Prnp*^{-/-} (Sakaguchi et al., 1996), ZH-II *Prnp*^{-/-} (Rossi et al., 2001), and Rcm0 mice (Moore et al., 1999), develop progressive cerebellar Purkinje cell degeneration with ataxia in advanced age. This phenotype was originally attributed to the lack of PrP^C and ran counter to the two PrP knockout mouse lines produced earlier: the ZH-I *Prnp*^{0/0} (Büeler et al., 1992) and the Edbg *Prnp*^{-/-} mice (Manson et al., 1994). The characterization of Nsgk *Prnp*^{-/-} mice was particularly conscientious: the authors reintroduced *Prnp* as a transgene by genetic crosses and showed that this manipulation rescued the Purkinje cell degeneration. It seemed entirely reasonable, hence, to conclude that PrP^C is necessary for cerebellar homeostasis. Yet this interpretation could not be easily reconciled with the lack of phenotype in the remaining knockout lines and eventually was proven to be incorrect.

The inconsistency was eventually resolved by David Westaway's discovery of a novel gene located just 16 kilobases downstream of *Prnp* and encoding a 179 residue protein that has sequence similarities to the C terminus of PrP and was thus termed Doppel or Dpl (Moore et al., 1999). It then emerged that the gene targeting strategy in all ataxic PrP-deficient mice was associated with deletion of a splice acceptor site located on the coding exon of *Prnp*. This modification effectively places Dpl under transcriptional control of the *Prnp* promoter. As a consequence, brain expression of Dpl, which is normally very low, skyrockets in Nsgk, ZH-II, and Rcm0 mice (Weissmann and Aguzzi, 1999). This is clearly neurotoxic, as ablation of the Dpl reading frame from ZH-II

mice abolishes the Purkinje cell degeneration phenotype (Nicolas Genoud, Axel Behrens, and A.A., unpublished data).

Most intriguingly, Dpl-dependent neurodegeneration is abolished by cell-autonomous coexpression of full-length PrP (Rossi et al., 2001). Formally, this indicates that Dpl and PrP^C act antagonistically, maybe because they bind to a hitherto conjectural common ligand (Figure 4A), which was provisionally termed L_{PrP} (Shmerling et al., 1998). Alternatively, PrP^C and Dpl might engage in heterooligomeric complexes (Figure 4B), whose function could depend on their stoichiometric composition (Behrens and Aguzzi, 2002). The same mechanism may be operative in transgenic mice produced by Doron Shmerling and Charles Weissmann (Shmerling et al., 1998) in an attempt to specify the domain of PrP^C required for prion replication. Expression of a PrP variant that lacks a large part of the N terminus of PrP in *Prnp*^{0/0} mice induces spontaneous cerebellar degeneration, which however affects granule cells rather than Purkinje cells (the promoter used was inactive in Purkinje cells) and can also be prevented by the coexpression of a single endogenous *Prnp* allele. Structural studies have shown that human Dpl contains a relatively short, flexibly disordered “tail” comprising residues 24-51 and a globular domain extending from residues 52 to 149 for which a detailed structure was obtained (Luhers et al., 2003). Despite their highly divergent primary sequence, Dpl is largely superimposable to the carboxy-proximal half of PrP^C.

The molecular pathways by which Dpl and amino-proximally truncated PrP damage the cerebellum are unknown. However, the suppressibility of both phenotypes by full-length PrP^C is indicative of a high degree of specificity. Therefore, we contend that this model presently represents the best validated window of entry to determine the function of PrP^C in vivo.

The Basis of Prion Neurotoxicity

PrP^{Sc} accumulation in the brain is the hallmark of prion diseases, and PrP^{Sc} is—for all we know—a major component of the infectious agent. But is PrP^{Sc} also directly responsible for the devastating CNS pathology typical of prion diseases? On the one hand, accumulation of amyloid (or preamyloid) in the CNS is likely to be generally unhealthy, as exemplified by Alzheimer's disease (Aguzzi and Haass, 2003) and cerebral vascular amyloidoses (Revesz et al., 2002). On the other hand, chronic deposition of PrP^{Sc} does not damage *Prnp* knockout brains (Brandner et al., 1996a), and depletion of PrP^C from neurons of scrapie-infected mice prevents disease (Mallucci et al., 2003). Therefore, accumulation of PrP^{Sc} is unlikely to fully account for prion pathology. If so, what is it that actually kills the neurons?

Brains of Creutzfeldt-Jakob disease victims look truly frightening. In heavily affected areas, there is hardly any neuron left, and the brain tissue texture is coarsened by the abnormal growth of astrocytes (“gliosis”) and microglial cells. The most telling hallmark is spongiosis, a peculiar microvacuolation affecting residual neural cells.

The molecular steps that emanate from prion replication and lead to such destruction are unknown. Some




Figure 4. Hypothetical Models for the Function of PrP^C and the Neurotoxicity of ΔPrP^C and Dpl
 (A) PrP^C and Dpl (or ΔPrP) may compete for a common ligand, provisionally termed L_{PrP}. In order to accommodate the lack of neurodegeneration in Prnp^{0/0} mice, however, one would have to postulate the existence of a functional PrP^C analog, here termed π. While this model accommodates all experimental findings known to date, no physical evidence has come forward for the existence of L_{PrP} and π.
 (B) Dpl and ΔPrP may form a homomultimeric toxic aggregate, which may be inactivated by participation of full-length PrP^C. Toxicity may come about by various hypothetical mechanisms. For example, if such aggregates were to span a membrane, toxic properties may relate to the formation of pores.

gain of toxic function is likely, as constitutive or postnatal depletion of PrP^C does not trigger any pathology. A lively discussion is developing on the role of abnormal PrP^C topologies. Targeting of PrP to the cytosol results in rapidly lethal neurodegeneration (yet without PrP^{Sc}), and proteasome inhibition induces a slightly protease-resistant PrP species in cultured cells, which may be self-sustaining—at least for a while (Ma and Lindquist, 2002; Ma et al., 2002). Therefore, prion toxicity may start with retrotranslocation of PrP^C from the endoplasmic reticulum to the cytosol, in conjunction with impaired proteasomal function. While PrP is clearly toxic in the cytosol, the details of how it may get there are debated. Cytosolic PrP retains its secretory leader peptide and does not contain a glycosyl phosphatidyl inositol anchor, suggesting that it never enters the endoplasmic reticulum (Drisaldi et al., 2003). Whether toxicity of cytosolic PrP is universal, however, is currently quite hotly discussed (Roucou et al., 2003). On the other hand, Lingappa found that PrP^C assumes a transmembrane topology (CtmPrP), whose concentration correlates with neurotoxicity (Hegde et al., 1998, 1999). These data suggest that CtmPrP represents a major toxic moiety.

We still know nothing of the biochemical pathways leading to brain damage, be they triggered by cytoplasmic PrP or by CtmPrP; these may lead to the identification of therapeutic targets and may share components with other neurodegenerative diseases.

The Future of Prion Therapeutics

An impressive wealth of molecules was touted as potential antiprion lead compounds. However, none of these therapeutic leads have proven their usefulness yet in clinical settings, and some have conspicuously failed. One of the possible problems derives from the fact that

most antiprion compounds were identified in cell culture assays, where chronically prion-infected neuroblastoma cells are “cured” of their PrP^{Sc} and prion burden. A startling variety of substances appears to possess such prion-curing properties; a nonexhaustive list includes compounds as diverse as Congo red (Caughey and Race, 1992), amphotericin B, anthracyclins (Tagliavini et al., 1997), sulfated polyanions (Caughey and Raymond, 1993), porphyrins (Priola et al., 2000), branched polyamines (Supattapone et al., 2001), “β sheet breakers” (Soto et al., 2000), and the spice curcumin (Caughey et al., 2003).

Disappointingly, none of these compounds proved very effective for actual therapy of sick animals—let alone patients. We therefore believe that it is premature to treat patients with alleged antiprion drugs on the sole basis of antiprion efficacy in neuroblastoma cells. This shortcut was taken in the case of quinacrine, which cures scrapie-infected cultured cells with impressive efficacy (Korth et al., 2001), yet appears to be utterly ineffective in scrapie-infected mice (Collins et al., 2002) and in CJD patients (Cooper, 2002), besides being severely hepatotoxic (Scoazec et al., 2003).

Why do scrapie-infected cells fare so poorly as a model system for prion therapy? In our experience, infection rarely hits all cells in any given culture, and the prion-infected state can be quite unstable. Therefore, one could speculate that a variety of stressors may masquerade as antiprion cures by conferring a selective advantage to noninfected cells. This interpretation would explain the puzzling observation that antiprion “cure” is brought about by compounds with no structural or biological similarities.

Cytidyl-guanyl oligodeoxynucleotides (CpG-ODN), which bind Toll-like receptor 9 (TLR9) and stimulate innate

immune responses, were reported to delay disease upon chronic administration to scrapie-infected mice (Sethi et al., 2002). The contention that immune stimulation might protect against prions is extraordinary and is difficult to reconcile with the observation that immune deficiencies of all kinds inhibit prion spread (Frigg et al., 1999; Klein et al., 1997, 1998, 2001; Prinz et al., 2003c). Besides, MyD88^{-/-} mice undergo normal prion pathogenesis despite abrogation of TLR9 signaling (Prinz et al., 2003b), and we could not evidence any major effects of TLR9 stimulation on the course of disease—in a paradigm identical to that described originally (M.P., M. Heikenwälder, and A.A., unpublished data). Instead, repeated CpG-ODN administration proved extremely lymphotoxic (Heikenwalder et al., 2004)—a fact that may well explain its antiprion properties.

On a more positive note, the tremendous interest in this field has attracted researchers from various neighboring disciplines, including immunology, genetics, and pharmacology, and therefore it is to hope that rational and efficient methods for managing prion infections will be developed in the future.

Immunotherapy against Prions?

Prions are sturdy and their resistance against sterilization is proverbial, yet exposure in vitro to anti-PrP antisera can reduce the titer of infectious hamster brain homogenates (Gabizon et al., 1988). Anti-PrP antibodies were found to inhibit formation of protease-resistant PrP in a cell-free system (Horiuchi and Caughey, 1999). Also, antibodies (Klein et al., 2001) and F(ab) fragments to PrP (Enari et al., 2001; Peretz et al., 2001) can suppress prion replication in cultured cells.

While these data suggest the feasibility of antiprion immunoprophylaxis, the mammalian immune system is essentially tolerant to PrP^C; this is hardly a surprise, given that PrP^C is expressed on T and B cells. Ablation of *Prnp* (Büeler et al., 1992) renders mice highly susceptible to immunization with prions (Brandner et al., 1996b), and indeed some of the best monoclonal antibodies to PrP^C were generated in *Prnp*^{0/0} mice (Prusiner et al., 1993).

Tolerance was circumvented by transgenic expression of an immunoglobulin μ chain containing the epitope-interacting region of 6H4, a high-affinity anti-PrP monoclonal antibody (Korth et al., 1997). The transgenic μ chain associated with endogenous κ and λ chains, some pairings lead to reactive moieties and, consequently, to high anti-PrP^C titers in *Prnp*^{0/0} and *Prnp*^{+/+} mice. The buildup of anti-PrP^C titers, however, was more sluggish in the presence of endogenous PrP^C, suggesting that clonal deletion is actually occurring. B cell clones with the highest affinity to PrP^C are probably eliminated by tolerance, while clones with medium affinity are retained (Figure 5A). The latter sufficed to block prion pathogenesis upon intraperitoneal prion inoculation (Heppner et al., 2001). Hence, B cells are not intrinsically tolerant to PrP^C and can, in principle, mount a protective humoral response against prions. It was then found, in a followup study, that passive transfer of anti-PrP monoclonal antibodies (in admittedly heroic amounts) can delay the onset of scrapie in mice infected with prions intraperitoneally, albeit not such infected intracerebrally (White et al., 2003).

The challenges to a practical antiprion immunization,

however, are enormous. While providing an encouraging proof of principle, transgenic immunization cannot easily be reduced to practice. Further, no protection was observed if treatment was started after the onset of clinical symptoms, suggesting that passive immunization might be a good candidate for prophylaxis rather than therapy of TSEs. Active immunization, like in most antiviral vaccines, may be more effective, but is rendered exceedingly difficult by the stringent tolerance to PrP^C (Souan et al., 2001; F. Heppner, E. Pelliccioli, M.P., and A.A., unpublished results; and Figure 5B).

Soluble Prion Antagonists

In several paradigms, expression of two PrP^C moieties subtly different from each other antagonizes prion replication. For example, humans heterozygous for a common *Prnp* polymorphism at codon 129 are largely protected from CJD: this effect is so important that it may have acted as selective evolutionary pressure (Mead et al., 2003). Similarly, transgenic expression of hamster PrP^C renders *Prnp*^{0/0} mice highly susceptible to hamster prions, whereas coexpression of mouse PrP^C diminishes this effect. Transdominant single nucleotide mutations of *Prnp* have also been described (Perrier et al., 2002).

The molecular basis for these effects is unknown; perhaps the subtly modified PrP^C acts as a decoy by binding incoming PrP^{Sc} (or protein X) and sequestering it into a complex incapable of further replication.

We tested the latter hypothesis by fusing an immunoglobulin Fc γ domain to PrP^C. The Fc γ tail served multiple purposes: (1) ligand dimerization, which may enhance its avidity for interacting partners; (2) provision of a convenient tag for affinity purification; (3) expression of the protein as a soluble moiety, which allows for testing cell-autonomous effects; and (4) increased stability in body fluids. Excitingly, the PrP-Fc γ fusion protein was found to compete with PrP^C for PrP^{Sc} (Figure 6) and to prolong the latency period of prion infection upon expression in transgenic mice (Meier et al., 2003). It will be exciting to determine whether PrP-Fc γ can act cell-autonomously when delivered as a drug. If that proves true, soluble prion protein mutants may represent useful prionostatic compounds.

Prion Diagnosis: Weaknesses and Challenges

Like in any other disease, early diagnosis would significantly advance the chances of success of any possible interventional approach. But when compared to other fields of microbiological diagnostics, the tools for prion diagnosis appear to be depressingly unsophisticated. Presymptomatic diagnosis is virtually impossible, and the earliest possible diagnosis is based on clinical signs and symptoms. Hence, prion infection is typically diagnosed after the disease has considerably progressed.

A significant advance in prion diagnostics was accomplished in 1997 by the discovery that protease-resistant PrP^{Sc} can be detected in tonsillar tissue of vCJD patients (Hill et al., 1997b). It was hence proposed that tonsil biopsy may be the method of choice for diagnosis of vCJD (Hill et al., 1999). Furthermore, there have been reports of individual cases showing detection of PrP^{Sc} at preclinical stages of the disease in tonsil (Schreuder et al., 1996) as well as in the appendix (Hilton et al., 1998), indicating that lymphoid tissue biopsy may be useful for




Figure 5. Affinity of Antibodies, Tolerance, and Immunity against Prions

(A) When forced to express a transgenic heavy chain with anti-PrP specificity, B lymphocytes may couple it to a large repertoire of endogenous light chains. Some of the VH-VL pairs (variable domains of heavy and light chains) may yield very high-affinity antibodies, whereas others will have low or no affinity. (B) Mendelian crosses of α -PrP-VH transgenic mice with *Prnp*^{0/0}, wild-type, and PrP^C-overexpressing transgenic mice informed on tolerogenic constraints. In the absence of endogenous PrP^C, mouse sera exhibited high anti-PrP^C titers. In wild-type mice, anti-PrP^C titers despite some clonal deletion, whereas massive overexpression of PrP^C led to dramatic lymphopenia (Heppner et al., 2001). Instead, active immunization yields consistently high anti-PrP^C titers only in *Prnp*^{0/0} mice. The permissivity of B lymphocytes to expression of anti-PrP^C specificities implies that tolerance to PrP^C is predominantly dictated by T-helper constraints. CH: Constant region of the heavy chain. Fab: antigen binding fragment. Fc μ : IgM-specific heavy chain.

diagnosing presymptomatic individuals. These observations triggered large screenings of human populations for subclinical vCJD prevalence using appendectomy and tonsillectomy specimens (Glatzel et al., 2003b). PrP^{Sc}-positive lymphoid tissue was long considered to

be a vCJD-specific feature that would not apply to any other forms of human prion diseases (Hill et al., 1999). However, a recent survey of peripheral tissues of patients with sporadic CJD has identified PrP^{Sc} in as many as one-third of skeletal muscle and spleen samples




Figure 6. A Model for the Antiprion Action of PrP-Fc₂

The template refolding model of prion replication (top) postulates a transient dimerization of PrP^C and PrP^{Sc}. As a result, PrP^{Sc} would impart its own β sheet-rich, protease-resistant conformation onto PrP^C. In the absence of PrP^C, soluble dimeric PrP does not support replication of the infectious agent, nor formation of a protease-resistant moiety (middle). Although several lines of evidence indicate that it can associate with PrP^{Sc}, this association is nonproductive. Mice coexpressing PrP^C and soluble dimeric PrP replicate prions and eventually develop scrapie. However, the kinetics with which scrapie pathology develops, prion infectivity replicates, and PrP^{Sc} accumulates is slower than in wild-type mice. All experimental evidence presented here suggests that PrP-Fc₂ sequesters incoming as well as nascent PrP^{Sc} and renders it unavailable for further template-directed conversion of PrP^C (bottom).

(Glatzel et al., 2003a), as well as the olfactory epithelium of patients suffering from sCJD (Zanusso et al., 2003). These unexpected findings raise the hope that minimally invasive diagnostic procedures may take the place of brain biopsy in intravital CJD diagnostics.

The sensitivity of PrP^{Sc} detection was significantly improved by the sodium phosphotungstic (NaPTA) precipitation method (Safar et al., 1998; Figure 1). By concentrating PrP^{Sc} prior to Western blot analysis, this procedure improves the sensitivity of diagnostic assays by as much as 4 orders of magnitude (Wadsworth et al., 2001). An interesting development was brought about by the conformation-dependent immunoassay (CDI), in which conformational differences of PrP isoforms are mapped by quantitating the relative binding of antibodies to denatured and native protein (Safar et al., 1998). Rather than relying on protease resistance, the CDI measures a variety of misfolded PrP isoforms, which may increase its sensitivity (Bellon et al., 2003; Safar et al., 2002).

Be this as it may, all techniques described above suffer from the fact that PrP^{Sc} continues to represent a surrogate marker for prion infectivity—since (1) PrP^{Sc} has not been incontrovertibly shown to be congruent with the prion, and (2) several manipulations in vitro and in vivo can render PrP^C protease resistant without bestowing infectivity on it (Jackson et al., 1999). Therefore, determination of prion infectivity by bioassay remains the golden standard; like in Pasteur's age, the concentration of the infectious agent is determined by inoculating serial dilutions of the test material into experimental animals, and the dilution at which 50% of the animals contract the disease (termed ID₅₀) is determined. Naturally, this system is riddled with inconveniences: scores of animals need to be sacrificed, and the incubation times are lengthy (transgenic overexpression of PrP^C can help, but only to some extent). Also, the method tends to be breathtakingly inaccurate: the inoculation schemes used in most studies typically suffer from standard errors of ± 1 order of magnitude!

A radical improvement of this situation is likely to be brought about by the use of prion-susceptible cell lines (Bosque and Prusiner, 2000; Race et al., 1987). The determination of prion infectivity endpoints in cultures of highly susceptible cells combines the sensitivity and intrinsic biological validity of the bioassay (i.e., direct measurement of the infectivity) with the speed and convenience of an in vitro methodology amenable to medium-throughput automation (Klohn et al., 2003).

Unresolved Problems in Prion Science

The study of prions has taken several unexpected directions over the past few years. However, the areas that are still obscure do not relate only to the details; some of them concern the core of the prion concept (Chesebro, 1998). In summary, there are five large groups of questions regarding the basic science of prion replication and of development of transmissible spongiform encephalopathies diseases:

- Which are the molecular mechanisms of prion replication? How does the disease-associated prion protein, PrP^{Sc}, achieve the conversion of its cellular sibling,

PrP^C, into a likeness of itself? Which other proteins assist this process? Can we inhibit this process? If so, how?

- What is the essence of prion strains, which are operationally defined as variants of the infectious agent capable of retaining stable phenotypic traits upon serial passage in syngeneic hosts? The existence of strains is very well known in virology, but it was not predicted to exist in the case of an agent that propagates epigenetically.
- How do prions reach the brain after having entered the body? Which molecules and which cell types are involved in this process of neuroinvasion? Which inhibitory strategies are likely to succeed?
- The mechanisms of neurodegeneration in spongiform encephalopathies is not understood. Which are the pathogenetic cascades that are activated upon accumulation of disease-associated prion protein and ultimately lead to brain damage?
- What is the physiological function of the highly conserved, normal prion protein, PrP^C? The *Prnp* gene encoding PrP^C was identified in 1985 (Basler et al., 1986; Oesch et al., 1985), *Prnp* knockout mice were described in 1992 (Büeler et al., 1992), and some PrP^C-interacting proteins have been identified (Oesch et al., 1990; Rieger et al., 1997; Yehiely et al., 2002; Zanata et al., 2002). Yet the function of PrP^C remains unknown!

The questions described above deserve to be addressed with a vigorous research effort. Their study is likely to yield fundamental insights into the characteristics of these novel and essentially mysterious agents and may yield useful leads for the diagnosis and therapy of prion diseases.

Acknowledgments

We thank Charles Weissmann and Susan Lindquist for suggestions and critical comments. The Aguzzi lab is supported by the University of Zurich and by grants of the Bundesamt für Bildung und Wissenschaft, the Swiss National Foundation, the NCCR on neural plasticity and repair, the US Department of Defense (National Prion Research Program), and the Stambach Foundation.

References

- Aguzzi, A. (1996). Between cows and monkeys. *Nature* 381, 734.
- Aguzzi, A., and Weissmann, C. (1996). Spongiform encephalopathies: a suspicious signature. *Nature* 383, 666–667.
- Aguzzi, A., and Weissmann, C. (1997). Prion research: the next frontiers. *Nature* 389, 795–798.
- Aguzzi, A., and Haass, C. (2003). Games played by rogue proteins in prion and Alzheimer's disease. *Science* 302, 814–818.
- Aguzzi, A., and Hardt, W.D. (2003). Dangerous liaisons between a microbe and the prion protein. *J. Exp. Med.* 198, 1–4.
- Alper, T., Cramp, W.A., Haig, D.A., and Clarke, M.C. (1967). Does the agent of scrapie replicate without nucleic acid? *Nature* 214, 764–766.
- Alper, T., Haig, D.A., and Clarke, M.C. (1966). The exceptionally small size of the scrapie agent. *Biochem. Biophys. Res. Commun.* 22, 278–284.
- Andrews, N.J., Farrington, C.P., Ward, H.J., Cousens, S.N., Smith, P.G., Molesworth, A.M., Knight, R.S., Ironside, J.W., and Will, R.G. (2003). Deaths from variant Creutzfeldt-Jakob disease in the UK. *Lancet* 361, 751–752.
- Basler, K., Oesch, B., Scott, M., Westaway, D., Walchli, M., Groth,

- D.F., McKinley, M.P., Prusiner, S.B., and Weissmann, C. (1986). Scrapie and cellular PrP isoforms are encoded by the same chromosomal gene. *Cell* 46, 417–428.
- Behrens, A., and Aguzzi, A. (2002). Small is not beautiful: antagonizing functions for the prion protein PrP(C) and its homologue Dpl. *Trends Neurosci.* 25, 150–154.
- Bellon, A., Seyfert-Brandt, W., Lang, W., Baron, H., Groner, A., and Vey, M. (2003). Improved conformation-dependent immunoassay: suitability for human prion detection with enhanced sensitivity. *J. Gen. Virol.* 84, 1921–1925.
- Bessen, R.A., Kocisko, D.A., Raymond, G.J., Nandan, S., Lansbury, P.T., and Caughey, B. (1995). Non-genetic propagation of strain-specific properties of scrapie prion protein. *Nature* 375, 698–700.
- Blättler, T., Brandner, S., Raeber, A.J., Klein, M.A., Voigtländer, T., Weissmann, C., and Aguzzi, A. (1997). PrP-expressing tissue required for transfer of scrapie infectivity from spleen to brain. *Nature* 389, 69–73.
- Bolton, D.C., McKinley, M.P., and Prusiner, S.B. (1982). Identification of a protein that purifies with the scrapie prion. *Science* 218, 1309–1311.
- Bosque, P.J., and Prusiner, S.B. (2000). Cultured cell sublines highly susceptible to prion infection. *J. Virol.* 74, 4377–4386.
- Brandner, S., Isenmann, S., Raeber, A., Fischer, M., Sailer, A., Kobayashi, Y., Marino, S., Weissmann, C., and Aguzzi, A. (1996a). Normal host prion protein necessary for scrapie-induced neurotoxicity. *Nature* 379, 339–343.
- Brandner, S., Raeber, A., Sailer, A., Blättler, T., Fischer, M., Weissmann, C., and Aguzzi, A. (1996b). Normal host prion protein (PrP^C) is required for scrapie spread within the central nervous system. *Proc. Natl. Acad. Sci. USA* 93, 13148–13151.
- Brown, D.R., Schulz-Schaeffer, W.J., Schmidt, B., and Kretzschmar, H.A. (1997). Prion protein-deficient cells show altered response to oxidative stress due to decreased SOD-1 activity. *Exp. Neurol.* 146, 104–112.
- Brown, D.R., Wong, B.S., Hafiz, F., Clive, C., Haswell, S.J., and Jones, I.M. (1999). Normal prion protein has an activity like that of superoxide dismutase. *Biochem. J.* 344, 1–5.
- Bruce, M.E., Will, R.G., Ironside, J.W., McConnell, I., Drummond, D., Suttie, A., McCordle, L., Chree, A., Hope, J., Birkett, C., et al. (1997). Transmissions to mice indicate that 'new variant' CJD is caused by the BSE agent. *Nature* 389, 498–501.
- Büeler, H.R., Fischer, M., Lang, Y., Bluethmann, H., Lipp, H.P., DeArmond, S.J., Prusiner, S.B., Aguet, M., and Weissmann, C. (1992). Normal development and behaviour of mice lacking the neuronal cell-surface PrP protein. *Nature* 356, 577–582.
- Büeler, H.R., Aguzzi, A., Sailer, A., Greiner, R.A., Autenried, P., Aguet, M., and Weissmann, C. (1993). Mice devoid of PrP are resistant to scrapie. *Cell* 73, 1339–1347.
- Cashman, N.R., Loertscher, R., Nalbantoglu, J., Shaw, I., Kascsak, R.J., Bolton, D.C., and Bendheim, P.E. (1990). Cellular isoform of the scrapie agent protein participates in lymphocyte activation. *Cell* 61, 185–192.
- Caughey, B., and Race, R.E. (1992). Potent inhibition of scrapie-associated PrP accumulation by congo red. *J. Neurochem.* 59, 768–771.
- Caughey, B., and Raymond, G.J. (1993). Sulfated polyanion inhibition of scrapie-associated PrP accumulation in cultured cells. *J. Virol.* 67, 643–650.
- Caughey, B., Raymond, L.D., Raymond, G.J., Maxson, L., Silveira, J., and Baron, G.S. (2003). Inhibition of protease-resistant prion protein accumulation in vitro by curcumin. *J. Virol.* 77, 5499–5502.
- Caughey, B.W., Dong, A., Bhat, K.S., Ernst, D., Hayes, S.F., and Caughey, W.S. (1991). Secondary structure analysis of the scrapie-associated protein PrP 27–30 in water by infrared spectroscopy. *Biochemistry* 30, 7672–7680.
- Chandler, R.L. (1961). Encephalopathy in mice produced by inoculation with scrapie brain material. *Lancet* 1, 1378–1379.
- Chesebro, B. (1998). BSE and prions: uncertainties about the agent. *Science* 279, 42–43.
- Chesebro, B., Race, R., Wehrly, K., Nishio, J., Bloom, M., Lechner, D., Bergstrom, S., Robbins, K., Mayer, L., and Keith, J.M. (1985). Identification of scrapie prion protein-specific mRNA in scrapie-infected and uninfected brain. *Nature* 315, 331–333.
- Collinge, J., Whittington, M.A., Sidle, K.C., Smith, C.J., Palmer, M.S., Clarke, A.R., and Jefferys, J.G. (1994). Prion protein is necessary for normal synaptic function. *Nature* 370, 295–297.
- Collinge, J., Sidle, K.C., Meads, J., Ironside, J., and Hill, A.F. (1996). Molecular analysis of prion strain variation and the aetiology of 'new variant' CJD. *Nature* 383, 685–690.
- Collins, S.J., Lewis, V., Brazier, M., Hill, A.F., Fletcher, A., and Masters, C.L. (2002). Quinacrine does not prolong survival in a murine Creutzfeldt-Jakob disease model. *Ann. Neurol.* 52, 503–506.
- Cooper, E. (2002). Quinacrine in possible or probable CJD: it is blinded investigators, not patients, who must be in equipoise over treatment. *BMJ* 324, 239.
- Creutzfeldt, H.G. (1920). Über eine eigenartige herdförmige Erkrankung des Zentralnervensystems. *Z ges Neurol Psychiatr* 57, 1–19.
- Cuille, J., and Chelle, P.L. (1939). Experimental transmission of trembling to the goat. *Comptes Rendus des Seances de l'Academie des Sciences* 208, 1058–1160.
- DeArmond, S.J., and Prusiner, S.B. (1995). Etiology and pathogenesis of prion diseases. *Am. J. Pathol.* 146, 785–811.
- Dickinson, A.G., Meikle, V.M., and Fraser, H. (1968). Identification of a gene which controls the incubation period of some strains of scrapie agent in mice. *J. Comp. Pathol.* 78, 293–299.
- Driscaldi, B., Stewart, R.S., Adles, C., Stewart, L.R., Quaglio, E., Biasini, E., Fioriti, L., Chiesa, R., and Harris, D.A. (2003). Mutant PrP is delayed in its exit from the endoplasmic reticulum, but neither wild-type nor mutant PrP undergoes retrotranslocation prior to proteasomal degradation. *J. Biol. Chem.* 278, 21732–21743.
- Duffy, P., Wolf, J., Collins, G., DeVoe, A.G., Streeten, B., and Cowen, D. (1974). Possible person-to-person transmission of Creutzfeldt-Jakob disease. *N. Engl. J. Med.* 290, 692–693.
- Enari, M., Flechsig, E., and Weissmann, C. (2001). Scrapie prion protein accumulation by scrapie-infected neuroblastoma cells abrogated by exposure to a prion protein antibody. *Proc. Natl. Acad. Sci. USA* 98, 9295–9299.
- Fischer, M.B., Roeckl, C., Parizek, P., Schwarz, H.P., and Aguzzi, A. (2000). Binding of disease-associated prion protein to plasminogen. *Nature* 408, 479–483.
- Frigg, R., Klein, M.A., Hegyi, I., Zinkernagel, R.M., and Aguzzi, A. (1999). Scrapie pathogenesis in subclinically infected B-cell-deficient mice. *J. Virol.* 73, 9584–9588.
- Gabizon, R., McKinley, M.P., Groth, D., and Prusiner, S.B. (1988). Immunoaffinity purification and neutralization of scrapie prion infectivity. *Proc. Natl. Acad. Sci. USA* 85, 6617–6621.
- Gajdusek, D.C., and Zigas, V. (1957). Degenerative disease of the central nervous system in New Guinea - the endemic occurrence of 'kuru' in the native population. *N. Engl. J. Med.* 257, 974–978.
- Gajdusek, D.C., Gibbs, C.J., and Alpers, M. (1966). Experimental transmission of a Kuru-like syndrome to chimpanzees. *Nature* 209, 794–796.
- Gajdusek, D.C., Gibbs, C.J., Jr., and Alpers, M. (1967). Transmission and passage of experimental "kuru" to chimpanzees. *Science* 155, 212–214.
- Gibbs, C.J., Jr., Gajdusek, D.C., Asher, D.M., Alpers, M.P., Beck, E., Daniel, P.M., and Matthews, W.B. (1968). Creutzfeldt-Jakob disease (spongiform encephalopathy): transmission to the chimpanzee. *Science* 161, 388–389.
- Glatzel, M., Heppner, F.L., Albers, K.M., and Aguzzi, A. (2001). Sympathetic innervation of lymphoreticular organs is rate limiting for prion neuroinvasion. *Neuron* 31, 25–34.
- Glatzel, M., Abela, E., Maissen, M., and Aguzzi, A. (2003a). Extraneural pathologic prion protein in sporadic Creutzfeldt-Jakob disease. *N. Engl. J. Med.* 349, 1812–1820.
- Glatzel, M., Ott, P.M., Lindner, T., Gebbers, J.O., Gmur, A., Wuest, W., Huber, G., Moch, H., Podvinec, M., Stamm, B., and Aguzzi, A.

- (2003b). Human prion diseases: epidemiology and integrated risk assessment. *Lancet Neurol.* 2, 757–763.
- Glatzel, M., Rogivue, C., Ghani, A., Streffer, J.R., Amsler, L., and Aguzzi, A. (2002). Incidence of Creutzfeldt-Jakob disease in Switzerland. *Lancet* 360, 139–141.
- Glockshuber, R., Hornemann, S., Billeter, M., Riek, R., Wider, G., and Wuthrich, K. (1998). Prion protein structural features indicate possible relations to signal peptidases. *FEBS Lett.* 426, 291–296.
- Gorodinsky, A., and Harris, D.A. (1995). Glycolipid-anchored proteins in neuroblastoma cells form detergent-resistant complexes without caveolin. *J. Cell Biol.* 129, 619–627.
- Griffith, J.S. (1967). Self-replication and scrapie. *Nature* 215, 1043–1044.
- Hadlow, W.J. (1959). Scrapie and kuru. *Lancet* 2, 289–290.
- Harmey, J.H., Doyle, D., Brown, V., and Rogers, M.S. (1995). The cellular isoform of the prion protein, PrP^c, is associated with caveolae in mouse neuroblastoma (N2a) cells. *Biochem. Biophys. Res. Commun.* 210, 753–759.
- Hegde, R.S., Mastrianni, J.A., Scott, M.R., DeFea, K.A., Tremblay, P., Torchia, M., DeArmond, S.J., Prusiner, S.B., and Lingappa, V.R. (1998). A transmembrane form of the prion protein in neurodegenerative disease. *Science* 279, 827–834.
- Hegde, R.S., Tremblay, P., Groth, D., DeArmond, S.J., Prusiner, S.B., and Lingappa, V.R. (1999). Transmissible and genetic prion diseases share a common pathway of neurodegeneration. *Nature* 402, 822–826.
- Heikenwalder, M., Polymenidou, M., Junt, T., Sigurdson, C., Wagner, H., Akira, S., Zinkernagel, R., and Aguzzi, A. (2004). Lymphoid follicle destruction and immunosuppression after repeated CpG oligodeoxynucleotide administration. *Nat. Med.* 10, 187–192.
- Heppner, F.L., Musahl, C., Arrighi, I., Klein, M.A., Rulicke, T., Oesch, B., Zinkernagel, R.M., Kalinke, U., and Aguzzi, A. (2001). Prevention of scrapie pathogenesis by transgenic expression of anti-prion protein antibodies. *Science* 294, 178–182.
- Hill, A.F., Butterworth, R.J., Joiner, S., Jackson, G., Rossor, M.N., Thomas, D.J., Frosh, A., Tolley, N., Bell, J.E., Spencer, M., et al. (1999). Investigation of variant Creutzfeldt-Jakob disease and other human prion diseases with tonsil biopsy samples. *Lancet* 353, 183–189.
- Hill, A.F., Desbruslais, M., Joiner, S., Sidle, K.C., Gowland, I., Collinge, J., Doey, L.J., and Lantos, P. (1997a). The same prion strain causes vCJD and BSE [letter]. *Nature* 389, 448–450.
- Hill, A.F., Zeidler, M., Ironside, J., and Collinge, J. (1997b). Diagnosis of new variant Creutzfeldt-Jakob disease by tonsil biopsy. *Lancet* 349, 99.
- Hilton, D.A., Fathers, E., Edwards, P., Ironside, J.W., and Zajicek, J. (1998). Prion immunoreactivity in appendix before clinical onset of variant Creutzfeldt-Jakob disease. *Lancet* 352, 703–704.
- Horiuchi, M., and Caughey, B. (1999). Specific binding of normal prion protein to the scrapie form via a localized domain initiates its conversion to the protease-resistant state. *EMBO J.* 18, 3193–3203.
- Horiuchi, M., Priola, S.A., Chabry, J., and Caughey, B. (2000). Interactions between heterologous forms of prion protein: binding, inhibition of conversion, and species barriers. *Proc. Natl. Acad. Sci. USA* 97, 5836–5841.
- Houston, F., Foster, J.D., Chong, A., Hunter, N., and Bostock, C.J. (2000). Transmission of BSE by blood transfusion in sheep. *Lancet* 356, 999–1000.
- Hsiao, K., Baker, H.F., Crow, T.J., Poulter, M., Owen, F., Terwilliger, J.D., Westaway, D., Ott, J., and Prusiner, S.B. (1989). Linkage of a prion protein missense variant to Gerstmann-Straussler syndrome. *Nature* 338, 342–345.
- Hutter, G., Heppner, F.L., and Aguzzi, A. (2003). No superoxide dismutase activity of cellular prion protein in vivo. *Biol. Chem.* 384, 1279–1285.
- Jackson, G.S., Hosszu, L.L., Power, A., Hill, A.F., Kenney, J., Saibil, H., Craven, C.J., Waltho, J.P., Clarke, A.R., and Collinge, J. (1999). Reversible conversion of monomeric human prion protein between native and fibrillogenic conformations. *Science* 283, 1935–1937.
- Jakob, A. (1921). Über eigenartige Erkrankungen des Zentralnervensystems mit bemerkenswertem anatomischem Befunde. (Spastische Pseudosklerose-Encephalomyelopathie mit disseminierten Degenerationsherden). *Z. ges. Neurol. Psychiatr.* 64, 147–228.
- Jarrett, J.T., and Lansbury, P.T., Jr. (1993). Seeding “one-dimensional crystallization” of amyloid: a pathogenic mechanism in Alzheimer’s disease and scrapie? *Cell* 73, 1055–1058.
- Kitamoto, T., Muramoto, T., Mohri, S., Doh-ura, K., and Tateishi, J. (1991). Abnormal isoform of prion protein accumulates in follicular dendritic cells in mice with Creutzfeldt-Jakob disease. *J. Virol.* 65, 6292–6295.
- Klein, M.A., Frigg, R., Flechsig, E., Raeber, A.J., Kalinke, U., Bluethmann, H., Bootz, F., Suter, M., Zinkernagel, R.M., and Aguzzi, A. (1997). A crucial role for B cells in neuroinvasive scrapie. *Nature* 390, 687–690.
- Klein, M.A., Frigg, R., Raeber, A.J., Flechsig, E., Hegyi, I., Zinkernagel, R.M., Weissmann, C., and Aguzzi, A. (1998). PrP expression in B lymphocytes is not required for prion neuroinvasion. *Nat. Med.* 4, 1429–1433.
- Klein, M.A., Kaeser, P.S., Schwarz, P., Weyd, H., Xenarios, I., Zinkernagel, R.M., Carroll, M.C., Verbeek, J.S., Botto, M., Walport, M.J., et al. (2001). Complement facilitates early prion pathogenesis. *Nat. Med.* 7, 488–492.
- Klohn, P.C., Stoltze, L., Flechsig, E., Enari, M., and Weissmann, C. (2003). A quantitative, highly sensitive cell-based infectivity assay for mouse scrapie prions. *Proc. Natl. Acad. Sci. USA* 100, 11666–11671.
- Knaus, K.J., Morillas, M., Swietnicki, W., Malone, M., Surewicz, W.K., and Yee, V.C. (2001). Crystal structure of the human prion protein reveals a mechanism for oligomerization. *Nat. Struct. Biol.* 8, 770–774.
- Kocisko, D.A., Come, J.H., Priola, S.A., Chesebro, B., Raymond, G.J., Lansbury, P.T., and Caughey, B. (1994). Cell-free formation of protease-resistant prion protein. *Nature* 370, 471–474.
- Korth, C., Stierli, B., Streit, P., Moser, M., Schaller, O., Fischer, R., Schulz-Schaeffer, W., Kretzschmar, H., Raeber, A., Braun, U., et al. (1997). Prion (PrP^{Sc})-specific epitope defined by a monoclonal antibody. *Nature* 390, 74–77.
- Korth, C., May, B.C., Cohen, F.E., and Prusiner, S.B. (2001). Acridine and phenothiazine derivatives as pharmacotherapeutics for prion disease. *Proc. Natl. Acad. Sci. USA* 98, 9836–9841.
- Kurschner, C., Morgan, J.I., Yehiely, F., Bamborough, P., Da Costa, M., Perry, B.J., Thinakaran, G., Cohen, F.E., Carlson, G.A., and Prusiner, S.B. (1995). The cellular prion protein (PrP) selectively binds to Bcl-2 in the yeast two-hybrid system: identification of candidate proteins binding to prion protein. *Brain Res. Mol. Brain Res.* 30, 165–168.
- Lacroute, F. (1971). Non-Mendelian mutation allowing ureidosuccinic acid uptake in yeast. *J. Bacteriol.* 106, 519–522.
- Li, L., and Lindquist, S. (2000). Creating a protein-based element of inheritance. *Science* 287, 661–664.
- Luhers, T., Riek, R., Guntert, P., and Wuthrich, K. (2003). NMR structure of the human doppel protein. *J. Mol. Biol.* 326, 1549–1557.
- Ma, J., and Lindquist, S. (2002). Conversion of PrP to a self-perpetuating PrP^{Sc}-like conformation in the cytosol. *Science* 298, 1785–1788.
- Ma, J., Wollmann, R., and Lindquist, S. (2002). Neurotoxicity and neurodegeneration when PrP accumulates in the cytosol. *Science* 298, 1781–1785.
- Mabbott, N.A., Bruce, M.E., Botto, M., Walport, M.J., and Pepys, M.B. (2001). Temporary depletion of complement component C3 or genetic deficiency of C1q significantly delays onset of scrapie. *Nat. Med.* 7, 485–487.
- Mallucci, G.R., Ratte, S., Asante, E.A., Linehan, J., Gowland, I., Jefferys, J.G., and Collinge, J. (2002). Post-natal knockout of prion protein alters hippocampal CA1 properties, but does not result in neurodegeneration. *EMBO J.* 21, 202–210.
- Mallucci, G., Dickinson, A., Linehan, J., Klohn, P.C., Brandner, S., and Collinge, J. (2003). Depleting neuronal PrP in prion infection prevents disease and reverses spongiosis. *Science* 302, 871–874.

- Manson, J.C., Clarke, A.R., Hooper, M.L., Aitchison, L., McConnell, I., and Hope, J. (1994). 129/Ola mice carrying a null mutation in PrP that abolishes mRNA production are developmentally normal. *Mol. Neurobiol.* **8**, 121–127.
- McKinley, M.P., Bolton, D.C., and Prusiner, S.B. (1983). A protease-resistant protein is a structural component of the scrapie prion. *Cell* **35**, 57–62.
- Mead, S., Stumpf, M.P., Whitfield, J., Beck, J.A., Poulter, M., Campbell, T., Uphill, J.B., Goldstein, D., Alpers, M., Fisher, E.M., and Collinge, J. (2003). Balancing selection at the prion protein gene consistent with prehistoric kurulike epidemics. *Science* **300**, 640–643.
- Meier, P., Genoud, N., Prinz, M., Maissen, M., Rulicke, T., Zurbriggen, A., Raeber, A.J., and Aguzzi, A. (2003). Soluble dimeric prion protein binds PrP(Sc) in vivo and antagonizes prion disease. *Cell* **113**, 49–60.
- Montrasio, F., Frigg, R., Glatzel, M., Klein, M.A., Mackay, F., Aguzzi, A., and Weissmann, C. (2000). Impaired prion replication in spleens of mice lacking functional follicular dendritic cells. *Science* **288**, 1257–1259.
- Moore, R.C., Hope, J., McBride, P.A., McConnell, I., Selfridge, J., Melton, D.W., and Manson, J.C. (1998). Mice with gene targeted prion protein alterations show that Prnp, Sinc and Prni are congruent. *Nat. Genet.* **18**, 118–125.
- Moore, R.C., Lee, I.Y., Silverman, G.L., Harrison, P.M., Strome, R., Heinrich, C., Karunaratne, A., Pasternak, S.H., Chishti, M.A., Liang, Y., et al. (1999). Ataxia in prion protein (PrP)-deficient mice is associated with upregulation of the novel PrP-like protein doppel. *J. Mol. Biol.* **292**, 797–817.
- Mouillet-Richard, S., Ermonval, M., Chebassier, C., Laplanche, J.L., Lehmann, S., Launay, J.M., and Kellermann, O. (2000). Signal transduction through prion protein. *Science* **289**, 1925–1928.
- Naslavsky, N., Stein, R., Yanai, A., Friedlander, G., and Taraboulos, A. (1997). Characterization of detergent-insoluble complexes containing the cellular prion protein and its scrapie isoform. *J. Biol. Chem.* **272**, 6324–6331.
- Oesch, B., Westaway, D., Walchli, M., McKinley, M.P., Kent, S.B., Aebersold, R., Barry, R.A., Tempst, P., Teplow, D.B., Hood, L.E., and Weissmann, C. (1985). A cellular gene encodes scrapie PrP 27–30 protein. *Cell* **40**, 735–746.
- Oesch, B., Teplow, D.B., Stahl, N., Serban, D., Hood, L.E., and Prusiner, S.B. (1990). Identification of cellular proteins binding to the scrapie prion protein. *Biochemistry* **29**, 5848–5855.
- Pan, K.M., Baldwin, M., Nguyen, J., Gasset, M., Serban, A., Groth, D., Mehlhorn, I., Huang, Z., Fletterick, R.J., Cohen, F.E., et al. (1993). Conversion of alpha-helices into beta-sheets features in the formation of the scrapie prion proteins. *Proc. Natl. Acad. Sci. USA* **90**, 10962–10966.
- Paramithiotis, E., Pinard, M., Lawton, T., LaBoissiere, S., Leathers, V.L., Zou, W.Q., Estey, L.A., Lamontagne, J., Lehto, M.T., Kondejewski, L.H., et al. (2003). A prion protein epitope selective for the pathologically misfolded conformation. *Nat. Med.* **9**, 893–899.
- Pattison, I.H., and Millson, G.C. (1961). Scrapie produced experimentally in goats with special reference to the clinical syndrome. *J. Comp. Pathol.* **71**, 101–108.
- Peretz, D., Williamson, R.A., Kaneko, K., Vergara, J., Leclerc, E., Schmitt-Ulms, G., Mehlhorn, I.R., Legname, G., Wormald, M.R., Rudd, P.M., et al. (2001). Antibodies inhibit prion propagation and clear cell cultures of prion infectivity. *Nature* **412**, 739–743.
- Perrier, V., Kaneko, K., Safar, J., Vergara, J., Tremblay, P., DeArmond, S.J., Cohen, F.E., Prusiner, S.B., and Wallace, A.C. (2002). Dominant-negative inhibition of prion replication in transgenic mice. *Proc. Natl. Acad. Sci. USA* **99**, 13079–13084.
- Popper, K. (1991). Selections from the logic of scientific discovery. In *The Philosophy of Science*, R. Boyd, P. Gaspar, and J.D. Trout, eds. (Cambridge, MA: The MIT Press), pp. 100–119.
- Prinz, M., Heikenwalder, M., Junt, T., Schwarz, P., Glatzel, M., Heppner, F.L., Fu, Y.X., Lipp, M., and Aguzzi, A. (2003a). Positioning of follicular dendritic cells within the spleen controls prion neuroinvasion. *Nature* **425**, 957–962. Published online October 15, 2003. doi:10.1038/nature02072.
- Prinz, M., Heikenwalder, M., Schwarz, P., Takeda, K., Akira, S., and Aguzzi, A. (2003b). Prion pathogenesis in the absence of Toll-like receptor signalling. *EMBO Rep.* **4**, 195–199.
- Prinz, M., Huber, G., Macpherson, A.J., Heppner, F.L., Glatzel, M., Eugster, H.P., Wagner, N., and Aguzzi, A. (2003c). Oral prion infection requires normal numbers of peyer's patches but not of enteric lymphocytes. *Am. J. Pathol.* **162**, 1103–1111.
- Priola, S.A., Raines, A., and Caughey, W.S. (2000). Porphyrin and phthalocyanine antiscrapie compounds. *Science* **287**, 1503–1506.
- Prusiner, S.B. (1982). Novel proteinaceous infectious particles cause scrapie. *Science* **216**, 136–144.
- Prusiner, S.B., Hadlow, W.J., Eklund, C.M., and Race, R.E. (1977). Sedimentation properties of the scrapie agent. *Proc. Natl. Acad. Sci. USA* **74**, 4656–4660.
- Prusiner, S.B., Groth, D.F., Cochran, S.P., Masiarz, F.R., McKinley, M.P., and Martinez, H.M. (1980). Molecular properties, partial purification, and assay by incubation period measurements of the hamster scrapie agent. *Biochemistry* **19**, 4883–4891.
- Prusiner, S.B., Groth, D.F., McKinley, M.P., Cochran, S.P., Bowman, K.A., and Kasper, K.C. (1981). Thiocyanate and hydroxyl ions inactivate the scrapie agent. *Proc. Natl. Acad. Sci. USA* **78**, 4606–4610.
- Prusiner, S.B., Groth, D., Serban, A., Koehler, R., Foster, D., Torchia, M., Burton, D., Yang, S.L., and DeArmond, S.J. (1993). Ablation of the prion protein (PrP) gene in mice prevents scrapie and facilitates production of anti-PrP antibodies. *Proc. Natl. Acad. Sci. USA* **90**, 10608–10612.
- Race, R.E., Fadness, L.H., and Chesebro, B. (1987). Characterization of scrapie infection in mouse neuroblastoma cells. *J. Gen. Virol.* **68**, 1391–1399.
- Revesz, T., Holton, J.L., Lashley, T., Plant, G., Rostagno, A., Ghiso, J., and Frangione, B. (2002). Sporadic and familial cerebral amyloid angiopathies. *Brain Pathol.* **12**, 343–357.
- Rieger, R., Edenhofer, F., Lasmezas, C.I., and Weiss, S. (1997). The human 37-kDa laminin receptor precursor interacts with the prion protein in eukaryotic cells. *Nat. Med.* **3**, 1383–1388.
- Riek, R., Hornemann, S., Wider, G., Billeter, M., Glockshuber, R., and Wuthrich, K. (1996). Nmr structure of the mouse prion protein domain Prp(121–231). *Nature* **382**, 180–182.
- Riek, R., Hornemann, S., Wider, G., Glockshuber, R., and Wuthrich, K. (1997). NMR characterization of the full-length recombinant murine prion protein, mPrP(23–231). *FEBS Lett.* **413**, 282–288.
- Rossi, D., Cozzio, A., Flechsig, E., Klein, M.A., Aguzzi, A., and Weissmann, C. (2001). Onset of ataxia and Purkinje cell loss in PrP null mice inversely correlated with Dpl level in brain. *EMBO J.* **20**, 1–9.
- Roucou, X., Guo, Q., Zhang, Y., Goodyer, C.G., and LeBlanc, A.C. (2003). Cytosolic prion protein is not toxic and protects against Bax-mediated cell death in human primary neurons. *J. Biol. Chem.* **278**, 40877–40881.
- Saborio, G.P., Permanne, B., and Soto, C. (2001). Sensitive detection of pathological prion protein by cyclic amplification of protein misfolding. *Nature* **411**, 810–813.
- Safar, J., Wille, H., Itri, V., Groth, D., Serban, H., Torchia, M., Cohen, F.E., and Prusiner, S.B. (1998). Eight prion strains have PrP(Sc) molecules with different conformations. *Nat. Med.* **4**, 1157–1165.
- Safar, J.G., Scott, M., Monaghan, J., Deering, C., Didorenko, S., Vergara, J., Ball, H., Legname, G., Leclerc, E., Solfrosi, L., et al. (2002). Measuring prions causing bovine spongiform encephalopathy or chronic wasting disease by immunoassays and transgenic mice. *Nat. Biotechnol.* **20**, 1147–1150.
- Sailer, A., Büeler, H., Fischer, M., Aguzzi, A., and Weissmann, C. (1994). No propagation of prions in mice devoid of PrP. *Cell* **77**, 967–968.
- Sakaguchi, S., Katamine, S., Nishida, N., Moriuchi, R., Shigematsu, K., Sugimoto, T., Nakatani, A., Kataoka, Y., Houtani, T., Shirabe, S., et al. (1996). Loss of cerebellar purkinje cells in aged mice homozygous for a disrupted Prp gene. *Nature* **380**, 528–531.
- Sakudo, A., Lee, D.C., Saeki, K., Nakamura, Y., Inoue, K., Matsumoto, Y., Itohara, S., and Onodera, T. (2003). Impairment of superoxide dismutase activation by N-terminally truncated prion protein

- (PrP) in PrP-deficient neuronal cell line. *Biochem. Biophys. Res. Commun.* 308, 660–667.
- Schmitt-Ulms, G., Legname, G., Baldwin, M.A., Ball, H.L., Bradon, N., Bosque, P.J., Crossin, K.L., Edelman, G.M., DeArmond, S.J., Cohen, F.E., and Prusiner, S.B. (2001). Binding of neural cell adhesion molecules (N-CAMs) to the cellular prion protein. *J. Mol. Biol.* 314, 1209–1225.
- Schreuder, B.E.C., Vankeulen, L.J.M., Vromans, M.E.W., Langeveld, J.P.M., and Smits, M.A. (1996). Preclinical test for prion diseases. *Nature* 381, 563.
- Schwartz, M. (2003). How the cows turned mad (University of California Press).
- Scoazec, J.Y., Krolak-Salmon, P., Casez, O., Besson, G., Thobois, S., Kopp, N., Perret-Liaudet, A., and Streichenberger, N. (2003). Quinacrine-induced cytolytic hepatitis in sporadic Creutzfeldt-Jakob disease. *Ann. Neurol.* 53, 546–547.
- Scott, M., Foster, D., Mirenda, C., Serban, D., Coufal, F., Waelchli, M., Torchia, M., Groth, D., Carlson, G., DeArmond, S.J., et al. (1989). Transgenic mice expressing hamster prion protein produce species-specific scrapie infectivity and amyloid plaques. *Cell* 59, 847–857.
- Sethi, S., Lipford, G., Wagner, H., and Kretzschmar, H. (2002). Post-exposure prophylaxis against prion disease with a stimulator of innate immunity. *Lancet* 360, 229–230.
- Shmerling, D., Hegyi, I., Fischer, M., Blattler, T., Brandner, S., Gotz, J., Rulicke, T., Flechsig, E., Cozzio, A., von Mering, C., et al. (1998). Expression of amino-terminally truncated PrP in the mouse leading to ataxia and specific cerebellar lesions. *Cell* 93, 203–214.
- Soto, C., Kascsak, R.J., Saborio, G.P., Aucouturier, P., Wisniewski, T., Prelli, F., Kascsak, R., Mendez, E., Harris, D.A., Ironside, J., et al. (2000). Reversion of prion protein conformational changes by synthetic beta-sheet breaker peptides. *Lancet* 355, 192–197.
- Souan, L., Tal, Y., Felling, Y., Cohen, I.R., Taraboulos, A., and Mor, F. (2001). Modulation of proteinase-K resistant prion protein by prion peptide immunization. *Eur. J. Immunol.* 31, 2338–2346.
- Supattapone, S., Wille, H., Uyechi, L., Safar, J., Tremblay, P., Szoka, F.C., Cohen, F.E., Prusiner, S.B., and Scott, M.R. (2001). Branched polyamines cure prion-infected neuroblastoma cells. *J. Virol.* 75, 3453–3461.
- Tagliavini, F., McArthur, R.A., Canciani, B., Giaccone, G., Porro, M., Bugiani, M., Lievens, P.M., Bugiani, O., Peri, E., Dall'Ara, P., et al. (1997). Effectiveness of anthracycline against experimental prion disease in Syrian hamsters. *Science* 276, 1119–1122.
- Tobler, I., Gaus, S.E., Deboer, T., Achermann, P., Fischer, M., Rulicke, T., Moser, M., Oesch, B., McBride, P.A., and Manson, J.C. (1996). Altered circadian activity rhythms and sleep in mice devoid of prion protein. *Nature* 380, 639–642.
- True, H.L., and Lindquist, S.L. (2000). A yeast prion provides a mechanism for genetic variation and phenotypic diversity. *Nature* 407, 477–483.
- Valleron, A.J., Boelle, P.Y., Will, R., and Cesbron, J.Y. (2001). Estimation of epidemic size and incubation time based on age characteristics of vCJD in the United Kingdom. *Science* 294, 1726–1728.
- Wadsworth, J.D.F., Joiner, S., Hill, A.F., Campbell, T.A., Desbruslais, M., Luthert, P.J., and Collinge, J. (2001). Tissue distribution of protease resistant prion protein in variant CJD using a highly sensitive immuno-blotting assay. *Lancet* 358, 171–180.
- Waggoner, D.J., Drisaldi, B., Bartnikas, T.B., Casareno, R.L., Prohaska, J.R., Gitlin, J.D., and Harris, D.A. (2000). Brain copper content and cuproenzyme activity do not vary with prion protein Expression level. *J. Biol. Chem.* 275, 7455–7458.
- Watarai, M., Kim, S., Erdenebaatar, J., Makino, S., Horiuchi, M., Shirahata, T., Sakaguchi, S., and Katamine, S. (2003). Cellular prion protein promotes Brucella infection into macrophages. *J. Exp. Med.* 198, 5–17.
- Weissmann, C. (1991). Spongiform encephalopathies. The prion's progress. *Nature* 349, 569–571.
- Weissmann, C., and Aguzzi, A. (1999). Perspectives: neurobiology. PrP's double causes trouble. *Science* 286, 914–915.
- Wells, G.A., Scott, A.C., Johnson, C.T., Gunning, R.F., Hancock, R.D., Jeffrey, M., Dawson, M., and Bradley, R. (1987). A novel progressive spongiform encephalopathy in cattle. *Vet. Rec.* 121, 419–420.
- Westaway, D., Goodman, P.A., Mirenda, C.A., McKinley, M.P., Carlson, G.A., and Prusiner, S.B. (1987). Distinct prion proteins in short and long scrapie incubation period mice. *Cell* 51, 651–662.
- White, A.R., Enever, P., Tayebi, M., Mushens, R., Linehan, J., Brandner, S., Anstee, D., Collinge, J., and Hawke, S. (2003). Monoclonal antibodies inhibit prion replication and delay the development of prion disease. *Nature* 422, 80–83.
- Wickner, R.B. (1994). [URE3] as an altered URE2 protein: evidence for a prion analog in *Saccharomyces cerevisiae*. *Science* 264, 566–569.
- Will, R.G., Ironside, J.W., Zeidler, M., Cousens, S.N., Estibeiro, K., Alperovitch, A., Poser, S., Pocchiari, M., Hofman, A., and Smith, P.G. (1996). A new variant of Creutzfeldt-Jakob disease in the UK. *Lancet* 347, 921–925.
- Wille, H., Michelitsch, M.D., Guenebaut, V., Supattapone, S., Serban, A., Cohen, F.E., Agard, D.A., and Prusiner, S.B. (2002). Structural studies of the scrapie prion protein by electron crystallography. *Proc. Natl. Acad. Sci. USA* 99, 3563–3568.
- Williams, E.S., and Young, S. (1980). Chronic wasting disease of captive mule deer: a spongiform encephalopathy. *J. Wildl. Dis.* 16, 89–98.
- Yehiely, F., Bamborough, P., Costa, M.D., Perry, B.J., Thinakaran, G., Cohen, F.E., Carlson, G.A., and Prusiner, S.B. (2002). Identification of Candidate Proteins Binding to Prion Protein. Volume 3, Number 4 (1997), pages 339–355. *Neurobiol. Dis.* 10, 67–68.
- Zanata, S.M., Lopes, M.H., Mercadante, A.F., Hajj, G.N., Chiarini, L.B., Nomizo, R., Freitas, A.R., Cabral, A.L., Lee, K.S., Juliano, M.A., et al. (2002). Stress-inducible protein 1 is a cell surface ligand for cellular prion that triggers neuroprotection. *EMBO J.* 21, 3307–3316.
- Zanusso, G., Ferrari, S., Cardone, F., Zampieri, P., Gelati, M., Fiorini, M., Farinazzo, A., Gardiman, M., Cavallaro, T., Bentivoglio, M., et al. (2003). Detection of pathologic prion protein in the olfactory epithelium in sporadic Creutzfeldt-Jakob disease. *N. Engl. J. Med.* 348, 711–719.
- Zijlstra, M., Bix, M., Simister, N.E., Loring, J.M., Raulet, D.H., and Jaenisch, R. (1990). Beta 2-microglobulin deficient mice lack CD4–8+ cytolytic T cells. *Nature* 344, 742–746.