

What is the definition of sports-related concussion: a systematic review

Paul McCrory,¹ Nina Feddermann-Demont,^{2,3} Jiří Dvořák,^{3,4} J David Cassidy,^{5,6,7} Andrew McIntosh,^{8,9} Pieter E Vos,¹⁰ Ruben J Echemendia,^{11,12} Willem Meeuwisse,¹³ Alexander A Tarnutzer^{2,3}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2016-097393>).

For numbered affiliations see end of article.

Correspondence to

Dr Alexander A Tarnutzer, Department of Neurology, University Hospital Zurich, Frauenklinikstr. 26, Zurich CH-8091, Switzerland; atarnutzer@gmail.com

Accepted 2 March 2017

Published Online First

6 March 2017

ABSTRACT

Objectives Various definitions for concussion have been proposed, each having its strengths and weaknesses. We reviewed and compared current definitions and identified criteria necessary for an operational definition of sports-related concussion (SRC) in preparation of the 5th Concussion Consensus Conference (Berlin, Germany). We also assessed the role of biomechanical studies in informing an operational definition of SRC.

Design This is a systematic literature review.

Data sources Data sources include MEDLINE, Embase, Cumulative Index to Nursing and Allied Health Literature, Cochrane Central Register of Clinical Trials and SPORT Discus (accessed 14 September 2016).

Eligibility criteria for selecting studies Eligibility criteria were studies reporting (clinical) criteria for diagnosing SRC and studies containing SRC impact data.

Results Out of 1601 articles screened, 36 studies were included (2.2%), 14 reported on criteria for SRC definitions and 22 on biomechanical aspects of concussions. Six different operational definitions focusing on clinical findings and their dynamics were identified. Biomechanical studies were obtained almost exclusively on American football players. Angular and linear head accelerations linked to clinically confirmed concussions demonstrated considerable individual variation.

Summary/conclusions SRC is a traumatic brain injury that is defined as a complex pathophysiological process affecting the brain, induced by biomechanical forces with several common features that help define its nature. Limitations identified include that the current criteria for diagnosing SRC are clinically oriented and that there is no gold/standard to assess their diagnostic properties. A future, more valid definition of SRC would better identify concussed players by demonstrating high predictive positive/negative values. Currently, the use of helmet-based systems to study the biomechanics of SRC is limited to few collision sports. New approaches need to be developed to provide objective markers for SRC.

INTRODUCTION

Sports-related concussion (SRC) is a well-recognised clinical entity. However, its pathophysiological basis remains poorly understood. In the broadest clinical sense, concussion is often defined as representing the immediate and transient symptoms of a mild traumatic brain injury (mTBI). Such operational definitions, however, do not give any insights into the underlying processes through which the brain is

impaired, nor do they distinguish different grades of severity, nor reflect insights into the persistence of symptoms and/or abnormalities on specific investigational modalities.

Since the 1970s, clinicians and scientists have begun to distinguish SRC from other causes of concussion and mTBI, such as seen in motor vehicle crashes, etc. Although this seems like an arbitrary separation from other forms of TBI,^{1,2} it is largely driven by sporting bodies who see the need to have clear and practical guidelines to determine recovery and safe return to play for athletes suffering from SRC. In addition, sports participation can be viewed as research laboratories to study SRC and mTBI, given the detailed SRC phenotype data that are typically available in many sports.³ Having said that, it is critical to understand that the lessons derived from non-sporting mTBI research informs the understanding of SRC (and vice versa), and this arbitrary separation of sporting versus non-sporting TBI should not be viewed simply as a dichotomous or exclusive view of TBI.

Over the past 50 years, various definitions of concussion and mTBI have been proposed by individual authors, different research groups and international bodies. These definitions, however, do not necessarily concur with one another, although there are a number of common elements. The definitional differences make the understanding of injury epidemiology problematic and management challenging.⁴

This systematic review offers an overview of previous concussion definitions and on the role of biomechanical studies in this context. These were specific review questions given to this writing group by the Concussion in Sport Group (CISG). It aims at providing an updated operational definition of concussion and is part of a series of articles written in preparation of the 5th Concussion Consensus Conference held in October 2016 in Berlin, Germany.

HISTORICAL PERSPECTIVE

The clinical manifestations of concussion as a transient neurological syndrome due to head shaking (or ‘commotion’) without structural brain injury have been known since the 10th century AD, when the Persian physician, Rhazes, first defined the condition.^{5–7} Following pioneering experimental primate studies demonstrating the transient and functional nature of concussion, the term *acceleration concussion* was proposed as the generic descriptor that should be applied to all forms of traumatic brain injury.⁸ Implicit in this concept is that the term



CrossMark

To cite: McCrory P, Feddermann-Demont N, Dvořák J, et al. *Br J Sports Med* 2017;**51**:877–887.

concussion should be synonymous with *traumatic brain injury* of all severities. A variation on this view holds that concussion refers to the mechanism of injury and motion of the brain within the skull rather than any clinical symptoms or pathology.⁹

Dating back to the 1930s, numerous anecdotal concussion severity-grading scales have been published reflecting the variety of prevailing theories in existence at the time.^{10–11} By 2001, there were approximately 45 published scales, none of which had been scientifically validated.^{12–13} Although not defining concussion per se, these scales broadly attempted to separate arbitrary levels of concussion severity based on clinical symptoms that would, in turn, inform management and return to play advice. These scales have been reviewed in detail elsewhere; however, it was the view of the CISG in 2001 that these should be abandoned and clinical measures of assessment and recovery be utilised to determine safe return to play.¹²

WHAT DO BIOMECHANICS TELL US ABOUT THE DEFINITION OF SRC?

Studies examining biomechanics and head injury (including TBI) demonstrate broadly that impacts causing SRC are less severe than impacts causing cranial fractures, intracranial haemorrhages and diffuse axonal injury.^{14–18}

Biomechanical studies of concussion demonstrate both differential effects on brain regions depending on the impact force, site of impact and bony architecture of the skull, as well as considerable intrinsic variation in tolerance to head impacts, which will depend on energy status and previous concussion and could be speculated to have a genetic basis also.¹⁹

Recent studies on SRC have reported head impact exposure patterns for specific sports, for example, American football, ice hockey and Australian football. Those studies report head impact characteristics including frequency, head kinematics, head impact location and injury outcome. To quantify head impacts, studies have used helmet-based systems, mouthguard/headband/skin sensors as well as videometric studies. The measured or estimated head kinematics provide a starting point to understanding brain tissue loading. Studies have aimed at identifying concussion thresholds for head acceleration (linear and angular).¹⁸ Within these studies, the use of instrumented helmets has provided information on head impact exposures, although there remains some debate regarding the accuracy and precision of these head kinematic measurements.

Biomechanical modelling demonstrates that brain loading patterns (stresses and strains) are not uniformly distributed due to factors including the brain geometry, brain tissue properties, bony architecture of the internal skull and connective tissue, for example, falx cerebri and tentorium cerebelli. This means that certain anatomical areas will have greater or lesser physiological or biochemical disturbance depending on the interplay of the external forces, the head's overall kinematic responses and brain sensitivity, such that resultant clinical symptoms may vary from person to person.^{20–23}

To which extent such measurements of head kinematic responses, for example, linear and angular head acceleration, may influence a definition of concussion has not been determined, and a complete biomechanical understanding of these injuries and related modelling to predict SRC remains a work in progress.²⁴

AIMS OF THE REVIEW

The aim of this review was to perform a systematic review of the literature as per the CISG protocol²⁵ and specifically addresses the following two questions:

1. What are the critical clinical criteria for an operational definition of SRC?
2. Do the published biomechanical studies inform us about the definition of SRC?

As per the CISG process, the systematic review was presented at the 5th Concussion Consensus Conference in Berlin, Germany. The updated definition of SRC was developed from the literature review and informed by public discussions and was finalised by the scientific committee.

MATERIALS AND METHODS

The overall methodology of the CISG systematic review process has been outlined in a separate paper published accompanying this review.²⁵

Data sources and searches

A literature search (MEDLINE, Embase, Cumulative Index to Nursing and Allied Health Literature (CINAHL), Cochrane Central Register of Clinical Trials (CRCT) and SPORTDiscus) was performed (14 September 2016) to identify English-language articles reporting on clinical criteria of a definition of SRC (aim 1) or on original data of the biomechanics of SRC in humans (aim 2). The MEDLINE (OVID) search strategy was translated for each database and is reported in online supplementary file 1.

Note that for aim 1 we only included manuscripts that made a significant contribution to the diagnostic criteria for SRC, that is, modified existing diagnostic criteria or presented new criteria. Manuscripts that referred to, discussed or compared existing definitions of SRC were not eligible. Whereas for aim 1 consensus papers and reviews were also considered, original contributions containing data on players with clinically confirmed SRC were eligible only for aim 2. The role of biomarkers in concussion was addressed in a separate review in this issue and therefore is not further investigated here.

A manual search of reference lists from eligible articles was performed. We did not seek to identify research abstracts from meeting proceedings or unpublished studies. Studies with five or more participants were eligible for aim 2. This review complies with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.²⁶

Study selection

All identified articles were subject to title and abstract screening by two independent reviewers (AAT and NFD). Full-text screening was applied to all abstracts considered eligible by at least one reviewer. Articles were selected using predetermined criteria (for exclusion criteria, see online supplementary file 1). Discrepancies in selection status and reasons for exclusion were settled between the two reviewers by discussion and adjunction of a third reviewer (JD) if needed.

Data extraction and data synthesis

Data extraction was performed by AAT and confirmed by NFD. For aim 1, we extracted the key components of the concussion definition and how the definition was established. Data synthesis in these studies focused on the description of predefined key features (symptom onset and duration, mechanism, loss of consciousness (LOC), resolution of symptoms and neuroimaging), on the distinction between concussion and mTBI, and on the number of citations. In studies reporting on biomechanical aspects of SRC (aim 2), we extracted key aspects such as the investigated sports, recording devices used, acceleration values

resulting in a concussion and concussion location, and determined the distribution of mean linear and rotational acceleration values and the mean (± 1 SD) distribution of head impact location among studies.

Quality Assessment of Diagnostic Accuracy Studies-2 (QUADAS-2) assessment of included studies

For included studies reporting original data on the biomechanics of concussion (aim 2), the risk of bias and applicability concerns were assessed by one reviewer (AAT). A second reviewer (NFD) confirmed ratings, and disagreements were resolved by discussion. We opted for the QUADAS-2 tailored study criteria, as they are widely used and recommended for the assessment of diagnostic accuracy studies.²⁷ We did not restrict inclusion further based on QUADAS-2 results. Note that we did not apply QUADAS-2 to consensus/statement papers providing definitions

of concussion (aim 1), as this was not appropriate for this sort of publications.

The QUADAS-2 tool consists of four core domains (patient selection, index test, reference standard and flow/timing).²⁷ Risk of bias is assessed for all four domains, and applicability is assessed for the first three domains. Thus, seven items per study are assessed. For each item the risk of bias is identified as 'high', 'low' or 'unclear'.

RESULTS

We identified 1601 citations for title/abstract screening and 123 articles for full-text screening. Eventually, 36 (2.2%) studies were included for quantitative synthesis (figure 1). Among the 36 studies included, we identified 14 manuscripts defining the term concussion according to our selection criteria (aim 1), whereas 22 manuscripts reported on biomechanical aspects of SRC (aim 2).




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow chart.

Table 1 Overview of included studies providing definitions of concussion*

Authors, year published	Responsible organisation	Type of manuscript	Restricted to sport concussion (yes/no)	Age group	Concussion grading system (yes/no)	Remarks
Congress of Neurological Surgeons, 1966 ³⁷	Committee on Head Injury Nomenclature	Consensus statement	No	All ages	No	
Anon, 1997 ⁴⁶	American Academy of Neurology	Consensus statement	Yes	All ages	Yes (mild/moderate/severe)	
Giza <i>et al</i> , 2013 ³²	American Academy of Neurology	Consensus statement	Yes	All ages	No	Revision of 1997 AAN consensus ⁴⁶
Aubry <i>et al</i> , 2002 ¹²	CISG	Consensus statement	Yes	All ages	No	
McCroly <i>et al</i> , 2005 ²⁸	CISG	Consensus statement	Yes	All ages	No	Revision of the 2002 consensus ¹²
McCroly <i>et al</i> , 2009 ²⁹	CISG	Consensus statement	Yes	All ages	No	Revision of the 2005 CISG consensus ²⁸
McCroly <i>et al</i> , 2013 ³⁰	CISG	Consensus statement	Yes	All ages	No	Revision of the 2009 CISG consensus ²⁹
Guskiewicz <i>et al</i> , 2004 ³⁴	National Athletic Trainers' Association	Consensus statement	Yes	All ages	No	Existing grading systems are reviewed ^{46 47}
Broglio <i>et al</i> , 2014 ³³	National Athletic Trainers' Association	Consensus statement	Yes	All ages	No	Revision of 2004 NATA consensus ³⁴
No authors listed, 2006 ³⁵	Team Physician Consensus statement (AAFP, AAOS, ACSM, AMSSM, AOSSM, AOASM)	Consensus statement	Yes	All ages	No	
Herring <i>et al</i> , 2011 ³⁶	Team Physician Consensus statement (AAFP, AAOS, ACSM, AMSSM, AOSSM, AOASM)	Consensus statement	Yes	All ages	No	Revision of 2006 Team Physician Consensus ³⁵
Harmon <i>et al</i> , 2013 ³¹	American Medical Society for Sports Medicine	Consensus statement	Yes	All ages	No	
Berg <i>et al</i> , 2014 ³⁹	NA	Original research (retrospective)	Yes	All ages	No	Analysis of concussion definitions on popular concussion-related websites
Boutis <i>et al</i> , 2015 ³⁸	NA	Original research (prospective)	No	Children (5–18 years)	No	Implementation of CISG concussion definition in paediatric ED.

AAFP, American Academy of Family Physicians; AAN, American Academy of Neurology; AAOS, American Academy of Orthopaedic Surgeons; ACSM, American College of Sports Medicine; AMSSM, American Medical Society for Sports Medicine; AOSSM, American Orthopaedic Society for Sports Medicine; AOASM, American Osteopathic Academy of Sports Medicine; CISG, concussion in sports group; ED, emergency department; NA, not available; NATA, National Athletic Trainers' Association.

*Studies are presented according to the year of publication, with revised versions of statements from a given body following the first version (in descending order).

Aim 1: Studies reporting on concussion definitions

From the 14 manuscripts focusing on defining the term concussion (table 1), 12 studies from 6 different organisations (CISG, American Medical Society for Sports Medicine (AMSSM), National Athletic Trainers' Association (NATA), American Academy of Neurology (AAN), Team Physician Consensus Group (joint statement by the American Academy of Family Physicians, the American Academy of Orthopaedic Surgeons, the American College of Sports Medicine, the AMSSM, the American Orthopaedic Society for Sports Medicine, and the American Osteopathic Academy of Sports Medicine) and Committee on Head Injury Nomenclature From the Congress of Neurological Surgeons) provided consensus-based operational definitions of concussion.^{12 28–37} One study provided original data on the implementation of a structured concussion assessment in a paediatric emergency department,³⁸ and one study reviewed definitions of concussion as provided on websites.³⁹

Comparison of the different consensus-based operational definitions identified

Key aspects of the six definitions included are provided in table 2. Although all but one definition³⁷ were proposed or revised within the last 5 years and were restricted to SRC, they differed in the level of detail and the domains addressed.

Comparing the key elements characterising a concussion in the proposed consensus-based definitions showed both similarities and discrepancies for some items (table 2). Whereas in all

definitions loss of consciousness (LOC) was considered optional for making the diagnosis, onset (immediate/rapid/within minutes), duration, mechanism of impairment and resolution were addressed only by some of the definitions (table 2).

The assessment of specific domains was described in four definitions. Clinical symptoms typically included physical, cognitive, emotional and sleep disturbances, whereas examination included physical signs and neurocognitive testing (for cognitive and neurobehavioural alterations). A statement regarding typical findings on standard structural neuroimaging was provided only by two groups (CISG and Team Physician Consensus).

The terms 'concussion' and 'mTBI' were handled differently in these consensus statements.^{30–33 36} Whereas concussion was considered a subset of TBI (CISG)³⁰ or mTBI (AMSSM),³¹ concussion and mTBI were used synonymously by others (Team Physician Consensus statement, AAN consensus and NATA).^{32 33 36}

Aim 2: Studies reporting on biomechanical aspects of concussion

Twenty-two manuscripts reporting on biomechanical aspects of SRC met our inclusion criteria. From two studies, the results were reported in more than one paper. For each study we considered only the most recent publication^{40 41} and did not include the others.^{22 24 42} From the remaining 19 manuscripts, key aspects were extracted (tables 3 and 4).

Table 2 Key components for identifying a concussion—comparison of different definitions*

Organisation, year published	Definition of concussion	Domains assessed	Key features	Definition mTBI? Link concussion - TBI?	Notes
Concussion in sports group, ¹² 2002, 2005, 2009, 2013	'A complex pathophysiological process affecting the brain, induced by traumatic biomechanical forces' that 'may be caused by a direct blow to the head, face, neck or elsewhere in the body with an impulsive force transmitted to the head'. Common features include: ▶ Rapid onset of short-lived impairment of neurological function that resolves spontaneously ▶ May result in neuropathological changes, but the acute clinical symptoms largely reflect a functional disturbance rather than a structural injury ▶ Graded set of neurological syndromes that may or may not involve an LOC. Resolution of the clinical and cognitive features typically follows a sequential course. In some cases symptoms may be prolonged. ▶ Typically associated with grossly normal structural neuroimaging studies.	▶ Clinical symptoms (physical, cognitive, emotional) ▶ Physical signs ▶ Cognitive impairment ▶ Neurobehavioural features ▶ Sleep disturbances	<i>Onset:</i> rapid <i>Duration:</i> short-lived <i>Mechanism of impairment:</i> functional disturbance rather than structural injury <i>LOC:</i> may or may not <i>Resolution:</i> sequential <i>Neuroimaging:</i> no abnormalities on standard structural neuroimaging	No (concussion is considered as subset of TBI)	This definition also comments on typical neuroimaging features in concussion.
American Medical Society for Sports Medicine, ³¹ 2013	'A concussion is defined as a traumatically induced transient disturbance of brain function and is caused by a complex pathophysiological process. Concussions have also been referred to as mild traumatic brain injuries (mTBI). While all concussions are mTBIs, not all mTBIs are concussions. Concussions are a subset of mTBIs, on the less-severe end of the brain injury spectrum and are generally self-limited in duration and resolution'.	▶ Symptoms and signs (physical, cognitive, emotional, sleep)	<i>Onset:</i> not specified <i>Duration:</i> transient, self-limited <i>Mechanism of impairment:</i> functional disturbance <i>LOC:</i> may or may not <i>Resolution:</i> not specified <i>Neuroimaging:</i> not specified	No (concussion is considered as subset of mTBI)	
National Athletic Trainers' Association, ^{33 34} 2004, 2014	'Trauma-induced alteration in mental status that may or may not involve loss of consciousness'	▶ Not specified	<i>Onset:</i> not specified <i>Duration:</i> not specified <i>Mechanism of impairment:</i> not specified <i>LOC:</i> may or may not <i>Resolution:</i> not specified <i>Neuroimaging:</i> not specified	No (concussion and mTBI are used interchangeably)	
American Academy of Neurology, ^{32 46} 1997, 2013	'Concussion is a trauma-induced alteration in mental status that may or may not involve loss of consciousness. Confusion and amnesia are the hallmarks of concussion. The confusional episode and amnesia may occur immediately after the blow to the head or several minutes later'.	▶ Clinical symptoms (physical, cognitive, emotional) ▶ Physical signs ▶ Cognitive impairment ▶ Sleep disturbances	<i>Onset:</i> immediate or within mins <i>Duration:</i> not specified <i>Mechanism of impairment:</i> not specified <i>LOC:</i> may or may not <i>Resolution:</i> not specified <i>Neuroimaging:</i> not specified	No (concussion and mTBI are used interchangeably)	1997 definition was used as the 2013 revision provided a shortened definition only.
Team Physician Consensus statements (AAFP, AAOS, ACSM, AMSSM, AOSSM, AOASM), ^{35 36} 2006, 2011	'Concussion or mild traumatic brain injury (mTBI) is a pathophysiological process affecting the brain induced by direct or indirect biomechanical forces'. Common features include: ▶ Rapid onset of usually short-lived neurological impairment, which typically resolves spontaneously ▶ Acute clinical symptoms that usually reflect a functional disturbance rather than structural injury ▶ A range of clinical symptoms that may or may not involve LOC ▶ Routine neuroimaging studies are typically normal	▶ Clinical symptoms and signs (cognitive, somatic, affective, sleep disturbances)	<i>Onset:</i> rapid <i>Duration:</i> short-lived <i>Mechanism of impairment:</i> functional disturbance rather than structural injury <i>LOC:</i> may or may not <i>Resolution:</i> spontaneous <i>Neuroimaging:</i> no abnormalities on standard structural neuroimaging	Combined definition for mTBI and concussion	This definition also comments on typical neuroimaging features in concussion.
Congress of Neurological Surgeons, ³⁷ 1966	'a clinical syndrome characterized by the immediate and transient post-traumatic impairment of neural function such as alteration of consciousness, disturbance of vision or equilibrium due to mechanical forces'.	▶ Not specified	<i>Onset:</i> immediate <i>Duration:</i> transient <i>Mechanism of impairment:</i> not specified <i>LOC:</i> may or may not <i>Resolution:</i> not specified <i>Neuroimaging:</i> not specified	Not specified	

*In case of revised concussion definitions from the same group, the most recent version was considered if not stated otherwise.

AAFP, American Academy of Family Physicians; AAOS, American Academy of Orthopaedic Surgeons; ACSM, American College of Sports Medicine; AMSSM, American Medical Society for Sports Medicine; AOSSM, American Orthopaedic Society for Sports Medicine; AOASM, American Osteopathic Academy of Sports Medicine; LOC, loss of consciousness; mTBI, mild traumatic brain injury.

Risk of bias assessment (QUADAS-2)

Risk of bias and applicability concerns for patient selection were rated as 'high' in six studies as only a subset of identified concussions were included (online supplementary file 2). For index testing, all studies were considered 'high' risk of bias as the biomechanical data were analysed and interpreted with the knowledge of the results from the reference test. Eleven studies that were either directly funded by or had one/several co-author(s) who had a vested financial interest in the accelerometry system used were rated as 'high' risk for applicability concerns of the index test. Regarding the reference standard, we considered all 22 studies as 'unclear' risk of bias, as it remains unknown if all concussions were identified by the medical staff.

Key findings of included studies

To quantify head impacts, 17/19 studies used helmet-based systems. The Head Impact Telemetry System (HITS) was used in 15 studies and instrumented mouthguards with 6 degrees-of-freedom (DOF) sensors were used in 2 studies. Video analysis was applied to a study cohort of rugby and Australian rules football players,²¹ providing more detailed analysis on a subset of players in a follow-up study.⁴¹ Different combinations of parameters were assessed in these studies. The methods by which concussions were diagnosed remained unclear in five studies, and only five studies adhered to published concussion consensus definitions.

The measured mean peak linear accelerations in concussed players (data from 13/15 studies using the HITS) ranged between 43.0 and 145.0 g. In between these boundaries, most studies reported mean peak linear acceleration values of approximately 100 g. For rotational acceleration, the mean peak values (n=9 studies) in concussed players ranged between 3620 and 7230 rad/s² (table 4). Results from studies using instrumented mouthguards or video-based analyses demonstrated a similar range for peak linear accelerations, whereas peak rotational accelerations tended to be higher (table 4).

Eight HITS-based studies in American football players (and ice hockey players in one of those studies) reported the location of head impacts (n=308), with frontal impacts identified most frequently (n=151; 49%). In a single study on head impact location in Australian rules football and rugby based on video analysis,²¹ 69% of the impacts were located on the side of the head. In a follow-up study including non-injured athletes, the proportion of impacts to the temporal region was significantly greater (p=0.05) for concussion cases compared with no-injury cases, 60% and 23%, respectively.⁴¹ It is important to recognise that structural TBI or cranial fractures did not occur in these studies, and the vast majority of head impacts recorded with the HIT system did not result in concussion; for example, approximately 0.02% of the impacts in Broglio *et al*⁴³ were associated with a diagnosis of concussion.

In summary, the reported peak linear and rotational acceleration values in players with clinically confirmed concussion showed large inter-individual variability, and cut-off values were proposed in a single study only. Noteworthy, all studies had a high risk of bias and a high risk for applicability concerns for at least one item, warranting caution in interpreting these studies.

DISCUSSION

Our review identified six consensus-based definitions of SRC. Among the different organisations providing definitions, the consensus statements from the CISG were cited most frequently

(with 1376 citations in Google Scholar for the 2009 consensus statement), followed by the statements from AAN, NATA and AMSSM (all around 500–700 citations) (table 5). Because the 1966 Statement of the Congress of Neurological Surgeons was issued prior to the recent popularity of this topic, the number of citations for this document must be interpreted with caution.

On the basis of the highest number of citations, the consensus statements from the CISG can be considered the opinion leader, although it is acknowledged that earlier definitions largely predate the era of internet citations.^{12 28–30} These consensus-based definitions showed different levels of detail and weighted distinct domains differently. We demonstrated limitations even in those definitions that provided additional clarifying statements.^{30 36}

Recommendations for a future operational definition of SRC

On the basis of this systematic review and consensus discussions, it is recommended that the key elements of a future *clinical or operational definition of SRC* ideally include some or all of the following elements:

- ▶ Biomechanics—the injury is caused by quantified direct or indirect force(s) to the brain;
- ▶ Physiology—a defined physiological disruption of brain function;
- ▶ Clinical—a range of evolving clinical symptoms and signs including an alteration in cognitive functioning or mental state (eg, confusion, disorientation, slowed thinking) that may or may not involve transient LOC;
- ▶ Neuroimaging—a defined abnormality on advanced imaging platforms reflecting the underlying physiological abnormalities and clinical features;
- ▶ Fluid biomarkers and genetics—defined abnormalities reflecting the underlying physiological disruption or injury.

Despite many publications and definitional attempts, these considerations leave several issues unanswered; notably, does being dazed, seeing stars or feeling dizzy in the absence of altered mental state constitute either concussion or mTBI? The definition also does not explain how known concussion modifiers influence the clinical presentation.

Berlin consensus definition of SRCs

The 2016 Berlin Consensus Conference operational definition ('Berlin definition') is presented below. This definition is based on the systematic review of the literature presented here, as well as the consensus discussions during the plenary session by the conference participants and the expert panel.

Concussion is a traumatic brain injury induced by biomechanical forces. Several common features that may be used in clinically defining the nature of a concussive head injury include the following:

- ▶ Concussion may be caused either by a direct blow to the head, face, neck or elsewhere on the body with an impulsive force transmitted to the head.
- ▶ Concussion typically results in the rapid onset of short-lived impairment of neurological function that resolves spontaneously. However, in some cases, signs and symptoms evolve over a number of minutes to hours.
- ▶ Concussion may result in neuropathological changes, but the acute clinical signs and symptoms largely reflect a functional disturbance rather than a structural injury and, as such, no abnormality is seen on standard structural neuroimaging studies.
- ▶ Concussion results in a range of clinical signs and symptoms that may or may not involve LOC. Resolution of the clinical

Table 3 Overview of included original publications reporting on biomechanical aspects of concussion*

Authors, year published	Times cited (Google) [†]	Times cited (Web of Science) [†]	Times cited		Investigated sports	Level of play	Players (n), age, gender (F/M/B)	Recording system	Impacts/concussions	Parameters assessed	Concussion definition used	Diagnosed by	Notes
			Study design	Study design									
Duma <i>et al</i> , 2005 ⁴⁸	250	144	Prospective, observational	American football	College	38, NR, M	HITS	3312/1	Linear/rotational acc, Δ head vel. GSI, HIC	NR	NR		
Guskiewicz <i>et al</i> , 2007 ²⁰	248	145	Prospective, observational	American football	College	88, 20.2 \pm 1.8 years, M	HITS	1 04714/13	Linear/rotational acc	Own definition	Team physician		
Greenwald <i>et al</i> , 2008 ⁴⁹	245	158	Prospective, observational	American football	NCAA division I schools/high schools	449, NR, M	HITS	289 916/17	Linear/rotational acc, GSI, HIC	AAN	Medical staff (not specified)		
Rowson <i>et al</i> , 2012 ³⁰	163	85	Prospective, observational	American football	College	335, NR, M	HITS (n=314); custom 6-DOF sensors (n=21)	301 034/57	Rotational acc	CISG	Medical staff (not specified)		
Broglio <i>et al</i> , 2010 ⁵¹	139	79	Prospective, observational	American football	High school	78, 16.7 \pm 0.8 years, M	HITS	54247/13	Linear/rotational acc,	AAN	Certified AT, physician		
Schobel <i>et al</i> , 2007 ⁵²	136	79	Prospective, observational	American football	College/high school	56, NR, M	HITS	541 54/6	Linear acc	NR	NR		
McIntosh <i>et al</i> , 2000 ⁴¹	100	72	Retrospective	Rugby, Australian rules football	Professionals	100, NR, M	Video	100/100	Head impact dynamics	NR	NR		
Brolinson <i>et al</i> , 2006 ⁵³	92	NA	Prospective, observational	American football	College	52, NR, M	HITS	11 604/3	Linear acc	NR	Medical staff (not specified)		
McAllister <i>et al</i> , 2012 ⁵⁴	76	45	Prospective, observational	American football, ice hockey	College/high school	10, NR, M	HITS	10/10	Linear/rotational acc	Own definition	Certified AT, physician		
Funk <i>et al</i> , 2012 ⁵⁵	72	37	Prospective, observational	American football	College	98, NR, M	HITS	371 28/4	Linear acc, HIC	NR	AT, physician		
Duhaime <i>et al</i> , 2012 ⁵⁶	59	35	Prospective, observational	American football, ice hockey	College	450, NR, B	HITS	486 594/48§	Linear/rotational acc	NR	Certified AT, physician		
Broglio <i>et al</i> , 2011 ⁴³	45	27	Prospective, observational	American football	High school	95, NR, M	HITS	102 218/20	Linear/rotational acc, HITsp	AAN	Certified AT, physician		
Beckwith <i>et al</i> , 2013 ⁴⁰	39	24	Prospective, observational	American football	College/high school	1208, NR, M	HITS	161 732/105	Linear/rotational acc, Δ head vel. GSI, HIC	NR	AT, physician	This dataset has previously been reported by Beckwith <i>et al</i> 2013. ⁴² To avoid duplicity, this study has been removed from the analysis.	
Duma and Rowson, 2009 ⁵⁷	26	5	Prospective, observational	American football	College	NR, NR, M	HITS	71 300/6	Linear acc	NR	Team physician		

Continued

Table 3 Continued

Authors, year published	Times cited (Google) [†]	Times cited (Web of Science) [†]	Study design	Investigated sports	Level of play	Players (n), age, gender (F/M/B)	Recording system	Impacts/concussions	Parameters assessed	Concussion definition used	Diagnosed by	Notes
King <i>et al</i> , 2015 ³⁶	23	8	Prospective, observational	Rugby	Amateur	38, 2.2±4 years, M	Instrumented mouthguard (custom-built 6-DOF sensor)	20 689/2	Linear/rotational acc	NR	Medical staff (not specified)	
McIntosh <i>et al</i> , 2014 ⁴¹	19	4	Prospective, observational	Rugby and Australian rules football	Professionals	40, NR, M	Rigid body simulations, data from video	40/27	Linear/rotational acc	NR	NR	The 27 cases with concussion were taken from a larger sample previously published. ²¹ Preliminary analyses of the 27 cases were provided by Freche and McIntosh 2009 ²⁴ and Patton <i>et al</i> 2013. ²²
Hernandez <i>et al</i> , 2015 ⁵⁹	17	4	Prospective, observational	American football, boxing, MMA	College/professional	33, NR, B	Custom 6-DOF acc device	513/2	18 different parameters	NR	NR	
Wilcox <i>et al</i> , 2015 ⁶⁰	7	4	Prospective, observational	Ice hockey	College	58, NR, F	HITS	NR/9	Linear/rotational acc, HITsp	Own definition	Certified AT, physician	
Beckwith <i>et al</i> , 2009 ⁶¹	6	NA	Prospective, observational	American football	College/high school	52, 18.9±2.3 years, M	HITS	71 390/52	Linear/rotational GSI, HIC15	AAN	Medical staff (not specified)	While this study included 901 players, this manuscript only reports on the 52 players with concussion.

AAN, American Academy of Neurology; acc, acceleration; AT, athletic trainer; B, both gender; C1SG, Concussion in Sports Group; DOF, degrees-of-freedom; F, females; GSI, Gadd Severity Index; HIC, Head Injury Criterion; HITS, Head Impact Telemetry System; HITsp, Head Impact Technology severity profile; M, males; MMA, mixed martial arts; NA, not available; NCAA, National Collegiate Athletic Association; NR, not reported.

*Studies are presented according to the number of citations in Google Scholar (in descending order).

†Accessed on 19 October 2016 (google.com).

‡Service provided by Web of Science, Thomson Reuters. Accessed on 19 October 2016.

§Detailed numbers on the 48 concussions recorded: American football (n=40), women's ice hockey (n=7) and men's ice hockey (n=1).

Table 4 Key aspects of studies reporting on the biomechanics of concussion (only studies using HITS)

Authors, year published	Sports investigated	Impacts on days with concussion (mean±1 SD, range)	Mean peak values in concussed players (±1 SD, range)		Location of head impact (n)					
			Linear acc (g)	Rotational acc (rad/s ²)	Front	Top	Side	Back	All	
HITS technology used for data acquisition (n=15)										
Duma <i>et al</i> , 2005 ⁴⁸	American football	NR	81.0 (NR)	NR	0	0	1	0	1	
Guskiewicz <i>et al</i> , 2007 ²⁰	American football	NR	102.8±30.7 (60.51–168.71)	5312±3950 (NR)	NR	NR	NR	NR	13	
Greenwald <i>et al</i> , 2008 ⁴⁹	American football	NR	NR	NR	8	3	5	1	17	
Rowson <i>et al</i> , 2012 ⁵⁰	American football	NR	NR	5022±1791 (NR)	33 (including back)	17	7	Pooled with front	57	
Broglio <i>et al</i> , 2010 ⁵¹	American football	NR	105±18 (NR)	7230±1158 (NR)	8	2	2	1	13	
Schnebel <i>et al</i> , 2007 ⁵²	American football	34±24 (NR)	127.0±25.5 (81.9–145.7)	NR	NR	NR	NR	NR	6	
Brolinson <i>et al</i> , 2006 ⁵³	American football	NR	103.3±42.3 (NR)	NR	1	0	1	1	3	
McAllister <i>et al</i> , 2012 ⁵⁴	American football, ice hockey	NR	73.6±21.3 (NR)	5025±1226 (NR)	NR	NR	NR	NR	10	
Funk <i>et al</i> , 2012 ⁵⁵	American football	NR	145±35 (NR)	NR	NR	NR	NR	NR	4	
Duhaime <i>et al</i> , 2012 ⁵⁶	American football, ice hockey	19.5±15.7 (NR)	86.1±42.6 (16.5–177.9)*	3620±2166 (183–7589)*	13	9	5	8	35	
Broglio <i>et al</i> , 2011 ⁴³	American football	25.0±18.3 (NR)	93.6±27.5 (NR)	6402.6±17 530.9 (NR)	11	3	5	1	20	
Beckwith <i>et al</i> , 2013 ⁴⁰	American football	25.8±22.7 (1–108)	102.5±33.8 (29.3–205.3)	3977±2272 (183–10 484)	48	26	17	14	105	
Duma and Rowson, 2009 ⁵⁷	American football	NR	132.3±49.8 (NR)	NR	NR	NR	NR	NR	6	
Wilcox <i>et al</i> , 2015 ⁶⁰	Ice hockey	NR	43.0±11.5 (NR)	4030±1435 (NR)	NR	NR	NR	NR	9	
Beckwith <i>et al</i> , 2009 ⁶¹	American football	NR	107±31 (NR)	7079±3408 (NR)	28	12	11	4	55	
<i>All studies</i>					151 (49.0)	73 (23.7)	54 (17.5)	30 (9.7)	350	
Instrumented mouthguard with custom-built 6-DOF sensor used for data acquisition (n=2)										
King <i>et al</i> , 2015 ⁵⁸	Rugby	NR	74.9±28.2 (NR)	7627.5±3263.6 (NR)	NR	NR	NR	NR	2	
Hernandez <i>et al</i> , 2015 ⁵⁹	American football, boxing, MMA	NR	95.5±14.8 (85–106)	9565±3571 (7040–12 090)	NR	NR	NR	NR	2	
Video-based analyses (n=1)										
McIntosh <i>et al</i> , 2000 ²¹	Rugby, Australian rules football	NR	NR	NR	19 [†]	0	67	11	97 [‡]	
McIntosh <i>et al</i> , 2014 ⁴¹	Rugby, Australian rules football	NR	103.4±29.5 (NR)	7951±3562 (NR)	7 [§]	0	18	2	27 [¶]	

*Calculations based on those 31 concussions with an identified single impact leading to symptoms. For impact location, 35 concussions were considered.

†This includes nine concussions to the face.

‡Impact location was not reported in three players.

§This includes three concussions to the face.

¶This study provides a more detailed analysis of the biomechanical aspects in a subset of concussed players previously published.²¹ Note that the impact location has already been reported by the previous study.

Acc, acceleration; HITS, Head Impact Telemetry System; MMA, mixed martial arts; NR, not reported.

and cognitive features typically follows a sequential course.

However, in some cases symptoms may be prolonged.

The clinical signs and symptoms of concussion cannot be explained by drug, alcohol, medication use, other injuries (such as cervical injuries, peripheral vestibular dysfunction, etc) or other comorbidities (eg, psychological factors or coexisting medical conditions).

Limitations and outlook

Our systematic review may have a potential publication bias and language bias as only published, English-language articles were considered. It is important to note that even if a healthcare professional is 'skilled', what reproducible and valid criteria do they use to make this 'diagnosis' of SRC? Thus, the major challenges in concussion diagnosis are the following:

1. There is no gold or reference standard measure of concussion.

2. There is marked variability in the diagnostic criteria, with no studies examining the measurement properties of the definitional criteria.
3. Given 1 and 2, there is likely a large amount of information bias inherent in all concussion studies because there is no validated case definition. Thus, information bias would be present in studies of risk, prevention, prognosis, intervention and long-term sequelae. This likely explains some of the variability of findings in these areas.

The challenge is to validate an operational definition of SRC based on clinical criteria until a 'proven' gold standard is discovered and validated. This is highlighted as a major area of research for the reasons outlined above. If an objective gold standard becomes available, the clinical criteria could then be tested against it to provide diagnostic metrics such as positive and negative predictive values and likelihood ratios. Another approach in the absence of an objective gold standard is to use

Table 5 Number of citations for individual studies

Authors, year published	Responsible organisation	Times cited (Google)*	Times cited (Web of Science)†
McCrary <i>et al</i> , 2009 ²⁹	CISG	1376	203
McCrary <i>et al</i> , 2013 ³⁰	CISG	1158	468
McCrary <i>et al</i> , 2005 ²⁸	CISG	848	383
Aubry <i>et al</i> , 2002 ¹²	CISG	826	180
Harmon <i>et al</i> , 2013 ³¹	AMSSM	532	186
Guskiewicz <i>et al</i> , 2004 ³⁴	National Athletic Trainers' Association	513	269
Giza <i>et al</i> , 2013 ³²	American Academy of Neurology	374	192
Anon, 1997 ⁴⁶	American Academy of Neurology	352	379
Broglio <i>et al</i> , 2014 ³³	National Athletic Trainers' Association	140	66
Herring <i>et al</i> , 2011 ³⁶	Team Physician Consensus statement (AAFP, AAOS, ACSM, AMSSM, AOSSM, AOASM)	109	0
Boutis <i>et al</i> , 2015 ³⁸	NA	6	2
Herring <i>et al</i> , 2006 ³⁵	Team Physician Consensus statement (AAFP, AAOS, ACSM, AMSSM, AOSSM, AOASM)	4	1
Berg <i>et al</i> , 2014 ³⁹	NA	3	0
Congress of Neurological Surgeons, 1966 ³⁷	Committee on Head Injury Nomenclature	NA	39

AAFP, American Academy of Family Physicians; AAOS, American Academy of Orthopaedic Surgeons; ACSM, American College of Sports Medicine; AMSSM, American Medical Society for Sports Medicine; AOSSM, American Orthopaedic Society for Sports Medicine; AOASM, American Osteopathic Academy of Sports Medicine; CISG, Concussion in Sport Group; NA, not available.

* Accessed on 19 October 2016 (google.scholar.com).

† Service provided by Web of Science, Thomson Reuters. Accessed on 19 October 2016.

the prognostic utility of a concussion definition. If it accurately predicts outcome, it could be a useful measure to identify and triage cases. Such a prediction rule would be more useful than a diagnosis based on unknown criteria.⁴⁴

Currently, the investigation of biomechanical aspects of SRC focuses on high school/college collision sports where players wear helmets. In these studies peak linear acceleration values in concussed players averaged approximately 100 g. This is in good agreement with a recent systematic review.¹⁸ These impacts are less severe than head impacts resulting in structural head injury, for example, intracranial haemorrhages or contusions, diffuse axonal injury, and cranial fractures.^{15–17,45} Methodological limitations in these studies must be considered, such as helmet fit, not using consensus-based concussion definitions and not reporting whether the diagnosis was made by skilled medical personal or not. Furthermore, lack of duplicate risk of bias assessment in our systematic review should be considered as a limitation. Developments are occurring in mini-accelerometer devices that can be worn by athletes who do not wear a helmet. These devices offer opportunities for research and clinical practice. At present, there is no evidence that accelerometer devices or video-based observations of athletes can provide a 'diagnosis' of concussion. Therefore, at present, head acceleration data should not be used in isolation to remove a player from play or allow a player to keep playing following a hit.

CONCLUSIONS

With regard to both aims addressed in this systematic review, newer technological advances may give important insights into the underlying pathophysiology and ultimately provide a platform to develop a clear definition in the future. It can be anticipated that, in the future, head injury and concussion may eventually be defined by the severity of clinical signs, as well as genetic, epigenetic, metabolomic, proteomic, advanced imaging findings and blood/cerebrospinal-fluid biomarkers.

What is already known?

- ▶ Over the past 50 years, various definitions of sports-related concussion (SRC) and mild traumatic brain injury (mTBI) have been proposed by individual authors, different research groups and international bodies.
- ▶ Currently, there is no gold or reference standard for SRC diagnosis, and the diagnostic properties of the various definitions have not been studied.
- ▶ The different definitions of SRC do not concur with one another, and this variability has likely caused information or diagnostic misclassification bias in past studies on the management and prognosis of SRC.

What are the findings?

- ▶ Among currently available consensus-based definitions of SRC, the consensus statement from the Concussion in Sport Group can be considered the opinion leader based on citation statistics.
- ▶ These consensus-based definitions, however, showed different levels of detail and weighted distinct domains differently. Limitations even in those definitions that provided additional clarifying statements were demonstrated, including the overlap/distinction between SRC and mTBI.
- ▶ Angular and linear head accelerations linked to clinically confirmed SRC demonstrated considerable individual variation.
- ▶ The 2016 Berlin Consensus Conference operational definition ('Berlin definition') is presented.

Author affiliations

- ¹The Florey Institute of Neuroscience and Mental Health, Heidelberg, Victoria, Australia
- ²Department of Neurology, University Hospital Zurich and University of Zurich, Zurich, Switzerland
- ³Swiss Concussion Center, Schulthess Clinic, Zurich, Switzerland
- ⁴Department of Neurology, Schulthess Clinic, Zurich, Switzerland
- ⁵Division of Health Care and Outcomes Research, Krembil Research Institute, University Health Network, Toronto, Ontario, Canada
- ⁶Division of Epidemiology, Institute of Health Policy, Management and Evaluation, Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada
- ⁷Department of Sport Science and Clinical Biomechanics, Faculty of Health, University of Southern Denmark, Odense, Denmark
- ⁸Australian Collaboration for Research into Injury in Sport and its Prevention (ACRISP), Federation University Australia, Ballarat, Victoria, Australia
- ⁹Monash University Accident Research Centre, Monash University, Melbourne, Australia
- ¹⁰Department of Neurology, Slingeland Ziekenhuis, Doetinchem, The Netherlands
- ¹¹UOC Concussion Care Clinic, State College, Pennsylvania, USA
- ¹²Department of Psychology, University of Missouri - Kansas City, State College, Pennsylvania, USA
- ¹³Faculty of Kinesiology, Sport Injury Prevention Research Centre, University of Calgary, Calgary, Alberta, Canada

Contributors PM conceived the study, drafted and critically reviewed the manuscript, and approved the final version of the manuscript. NF-D performed the literature search, reviewed identified manuscripts and selected those matching inclusion criteria, edited and critically reviewed the manuscript, and approved the final version of the manuscript. JD conceived the study, edited and critically reviewed the manuscript, and approved the final version of the manuscript. J DC edited and critically reviewed the manuscript and approved the final version of the manuscript. AM edited and critically reviewed the manuscript and approved the final version of the manuscript. PEV edited and critically reviewed the manuscript and approved the final version of the manuscript. RJE edited and critically reviewed the manuscript and approved the final version of the manuscript. WM conceived the study, edited and critically reviewed the manuscript, and approved the final version of the manuscript. AAT performed the literature search, reviewed identified manuscripts and selected those matching inclusion criteria, performed the statistical analysis, drafted and critically reviewed the manuscript, and approved the final version of the manuscript.

Competing interests PM is a co-investigator, collaborator or consultant on grants relating to mild TBI funded by several governmental organisations. He is directly employed by the National Health and Medical Research Council of Australia and is based at the Florey Institute of Neuroscience and Mental Health. He is co-chair of the Australian Centre for Research into Sports Injury and its Prevention (ACRISP), which is one of the International University Research Centres for Prevention of Injury and Protection of Athlete Health supported by the International Olympic Committee (IOC). He is co-chair of the International Concussion in Sport Group. He has a clinical and consulting practice in general and sports neurology. He receives book royalties from McGraw-Hill and was employed in an editorial capacity by the British Medical Journal Publishing Group from 2001 to 2008. He has been reimbursed by the government, professional scientific bodies and sporting bodies for travel costs related to presenting research on mild TBI and sports-related concussion at meetings, scientific conferences and symposiums. He received consultancy fees in 2010 from Axon Sports (USA) for the development of educational material (which was not renewed) and has received research funding since 2001 from CogState Inc. He has not received any research funding, salary or other monies from the Australian Football League, FIFA or the National Football League. The Australian Football League funds research at the Florey Institute under a legal memorandum and PM does not receive any money from this industry-funded research. PM is a cofounder and shareholder in two biomedical companies (involved in eHealth and compression garment technologies) but does not hold any individual shares in any company related to concussion or brain injury assessment or technology. He did not receive any form of financial support directly related to this manuscript. NF-D and JD report no conflicts of interest. AM is a self-employed consultant and holds honorary academic appointments. He has been funded as a consultant to undertake research and policy related work for government, sports federations and industry on the topic of safety and injury risk management, including concussion, head injury and helmets. He did not receive any form of financial support directly related to this manuscript. He reports no conflicts of interest. PEV is a neurologist and Chairman of the Steering Committee of a trial in traumatic brain injury sponsored by EVER Neuro Pharma GmbH. He has been funded as a consultant for EVER Neuro Pharma. He did not receive any form of financial support directly related to this manuscript. He reports no conflicts of interest. JDC is an injury epidemiologist employed by the University Health Network, University of Toronto. He is a professor in the Division of Epidemiology and the Institute of Health Policy, Management and Evaluation at the Dalla Lana School of Public Health, University of Toronto. He has received funding from the Ontario Neurotrauma Foundation to undertake systematic reviews

on mild traumatic brain injury and served on the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury, which received funding from various sources, including traffic insurance companies in North America and Europe. He was retained as an expert witness for the National Hockey League on November 30, 2016. He did not receive any form of financial support related to this manuscript, and has no other conflicts of interest to report. RJE is a consultant to the National Hockey League, Major League Soccer, US Soccer Federation and Princeton University. He receives financial remuneration for these consulting relationships. He has a clinical practice in sport neuropsychology and serves as an expert (neuropsychology and sport neuropsychology) in medico-legal cases involving traumatic brain injury. WM is the Medical Director of the National Hockey League. He has received research grant support through the University of Calgary from the Canadian Institutes of Health Research, Alberta Innovates Health Solutions, the International Football Association (FIFA), Alberta Children's Hospital Research Institute and the Hotchkiss Brain Institute. He is the founding Chair of the Sport Injury Prevention Research Centre, which is one of the International Research Centres for Prevention of Injury and Protection of Athlete Health supported by the International Olympic Committee (IOC). He has a clinical consulting practice in sport medicine at the University of Calgary Sport Medicine Centre. He is an Expert Group member of the IOC Medical Commission and has received travel funding in that capacity.

Provenance and peer review Not commissioned; externally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Langlois JA, Rutland-Brown W, Wald MM. The epidemiology and impact of traumatic brain injury: a brief overview. *J Head Trauma Rehabil* 2006;21:375–8.
- Langlois JA, Sattin RW. Traumatic brain injury in the united states: research and programs of the centers for disease control and prevention (CDC). *J Head Trauma Rehabil* 2005;20:187–8.
- Kelly JP, Rosenberg JH. The development of guidelines for the management of concussion in sports. *J Head Trauma Rehabil* 1998;13:53–65.
- Ruff RM, Jurica P. In search of a unified definition for mild traumatic brain injury. *Brain Inj* 1999;13:943–52.
- Rhazes A. *Lutetiae ex officina*. Rome, Italy: Stephan, R, 1548.
- McCrorry PR, Berkovic SF. Concussion: the history of clinical and pathophysiological concepts and misconceptions. *Neurology* 2001;57:2283–9.
- Pearce JM. Observations on concussion. A review. *Eur Neurol* 2008;59(3-4):113–9.
- Denny-Brown D, Russell WR. Experimental cerebral concussion. *Brain* 1941;64:93–164.
- Russell WR, Smith A. Post-traumatic amnesia in closed head injury. *Arch Neurol* 1961;5:4–17.
- Russell WR. The after-effects of head injury. *Tr Med-Chir Soc Edinburgh* 1934;113:129–44.
- Russell W. Amnesia following head injuries* 1. *The Lancet* 1935;226:762–3.
- Aubry M, Cantu R, Dvorak J, et al; Concussion in Sport Group. Summary and agreement statement of the first international conference on concussion in sport, Vienna 2001. recommendations for the improvement of safety and health of Athletes who may suffer concussive injuries. *Br J Sports Med* 2002;36:6–10.
- Johnston KM, McCrorry P, Mohtadi NG, et al. Evidence-based review of sport-related concussion: clinical science. *Clin J Sport Med* 2001;11:150–9.
- Mertz HJ, ed. Head injury risk assessment for forehead impacts, SAE 960099. Proceedings of the SAE Technical Paper Series, 1996
- Prasad P, ed. The position of the U.S. delegation to the ISO Working Group 6 on the use of HIC in the automotive environment. *Proceedings of the SAE Technical Paper Series*, 1985.
- McIntosh A. Head and neck injury resulting from low velocity direct impact. Proceedings of the 37th STAPP Car Crash Conference, San Antonio, TX, 1993
- McIntosh AS. Biomechanical considerations in the design of equipment to prevent sports injury, Proceedings of the institution of mechanical engineers, part P. *J Sport Eng and Technol* 2012;226(3-4):193–9.
- Brennan JH, Mitra B, Synnot A, et al. Accelerometers for the assessment of concussion in male Athletes: a systematic review and Meta-Analysis. *Sports Med* 2017;47:469–78.
- Hayes JP, Bigler ED, Verfaellie M. Traumatic brain injury as a disorder of brain connectivity. *J Int Neuropsychol Soc* 2016;22:120–37.
- Guskiewicz KM, Mihalik JP, Shankar V, et al. Measurement of head impacts in collegiate football players: relationship between head impact biomechanics and acute clinical outcome after concussion. *Neurosurgery* 2007;61:1244–53.
- McIntosh AS, McCrorry P, Comerford J. The dynamics of concussive head impacts in rugby and Australian rules football. / Dynamique des chocs au niveau de la tete avec commotion cerebrale en rugby et en football australien. *Med Sci Sports Exerc* 2000;32:1980–4.
- Patton DA, McIntosh AS, Kleiven S. The biomechanical determinants of concussion: finite element simulations to investigate brain tissue deformations during sporting impacts to the unprotected head. *J Appl Biomech* 2013;29:721–30.

Review

- 23 Patton DA, McIntosh AS, Kleiven S. The biomechanical determinants of concussion: finite element simulations to investigate Tissue-Level predictors of injury during sporting impacts to the unprotected head. *J Appl Biomech* 2015;31:264–8.
- 24 Fréchède B, McIntosh AS. Numerical reconstruction of real-life concussive football impacts. *Med Sci Sports Exerc* 2009;41:390–8.
- 25 Meeuwisse W, Schneider K, Dvorak J, et al. The berlin 2016 process: a summary of methodology for the 5th international consensus conference on concussion in sport. *Br J Sports Med* (accepted and in press 22 Jan 2017).
- 26 Moher D, Liberati A, Tetzlaff J, et al; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- 27 Whiting PF, Rutjes AW, Westwood ME, et al; QUADAS-2 Group. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med* 2011;155:529–36.
- 28 McCrory P, Johnston K, Meeuwisse W, et al. Summary and agreement statement of the 2nd international conference on concussion in sport, Prague 2004. *Br J Sports Med* 2005;39:i78–86.
- 29 McCrory P, Meeuwisse W, Johnston K, et al. Consensus Statement on Concussion in Sport: the 3rd International Conference on Concussion in Sport held in Zurich, November 2008. *Br J Sports Med* 2009;43(Suppl 1):i76–84.
- 30 McCrory P, Meeuwisse WH, Aubry M, et al. Consensus statement on concussion in sport: the 4th International Conference on Concussion in Sport held in Zurich, November 2012. *Br J Sports Med* 2013;47:250–8.
- 31 Harmon KG, Drezner JA, Gammons M, et al. American Medical Society for Sports Medicine position statement: concussion in sport. *Br J Sports Med* 2013;47:15–26.
- 32 Giza CC, Kutcher JS, Ashwal S, et al. Summary of evidence-based guideline update: evaluation and management of concussion in sports. *Neurology* 2013;80:2250–7.
- 33 Broglio SP, Cantu RC, Gioia GA, et al; National Athletic Trainer's Association. National Athletic Trainers' Association position statement: management of sport concussion. *J Athl Train* 2014;49:245–65.
- 34 Guskiewicz KM, Bruce SL, Cantu RC, et al. National athletic trainers' Association position statement: management of Sport-Related concussion. *J Athl Train* 2004;39:280–97.
- 35 Anon. Concussion (mild traumatic brain injury) and the team physician: a consensus statement. *Med Sci Sports Exerc* 2006;38:395–9.
- 36 Herring SA, Cantu RC, Guskiewicz KM, et al; American College of Sports Medicine. Concussion (mild traumatic brain injury) and the team physician: a consensus statement--2011 update. *Med Sci Sports Exerc* 2011;43:2412–22.
- 37 Congress of neurological surgeons. Committee on head injury nomenclature: glossary of head injury. *Clin Neurosurg* 1966;12:386–94.
- 38 Boutis K, Weerdenburg K, Koo E, et al. The diagnosis of concussion in a pediatric emergency department. *J Pediatr* 2015;166:1214–20.
- 39 Berg GM, Hervey AM, Atterbury D, et al. Evaluating the quality of online information about concussions. *Jaapa* 2014;27:1–8.
- 40 Beckwith JG, Greenwald RM, Chu JJ, et al. Timing of Concussion diagnosis is related to head impact exposure prior to injury. *Med Sci Sports Exerc* 2013;45:747–54.
- 41 McIntosh AS, Patton DA, Fréchède B, et al. The biomechanics of concussion in unhelmeted football players in Australia: a case-control study. *BMJ Open* 2014;4:e005078.
- 42 Beckwith JG, Greenwald RM, Chu JJ, et al. Head impact exposure sustained by football players on days of diagnosed concussion. *Med Sci Sports Exerc* 2013;45:737–46.
- 43 Broglio SP, Eckner JT, Surma T, et al. Post-concussion cognitive declines and symptomatology are not related to concussion biomechanics in high school football players. *J Neurotrauma* 2011;28:2061–8.
- 44 Croft P, Altman DG, Deeks JJ, et al. The science of clinical practice: disease diagnosis or patient prognosis? Evidence about "what is likely to happen" should shape clinical practice. *BMC Med* 2015;13:20.
- 45 Margulies SS, Thibault LE. A proposed tolerance criterion for diffuse axonal injury in man. *J Biomech* 1992;25:917–23.
- 46 Anon. Practice parameter: the management of concussion in sports (summary statement). Report of the quality standards subcommittee. *Neurology* 1997;48:581–5.
- 47 Cantu RC. Posttraumatic retrograde and anterograde amnesia: pathophysiology and implications in grading and safe return to play. *J Athl Train* 2001;36:244–8.
- 48 Duma SM, Manoogian SJ, Bussone WR, et al. Analysis of real-time head accelerations in collegiate football players. *Clin J Sport Med* 2005;15:3–8.
- 49 Greenwald RM, Gwin JT, Chu JJ, et al. Head impact severity measures for evaluating mild traumatic brain injury risk exposure. *Neurosurgery* 2008;62:789–98.
- 50 Rowson S, Duma SM, Beckwith JG, et al. Rotational head kinematics in football impacts: an injury risk function for concussion. *Ann Biomed Eng* 2012;40:1–13.
- 51 Broglio SP, Schnebel B, Sosnoff JJ, et al. Biomechanical properties of concussions in high school football. *Med Sci Sports Exerc* 2010;42:2064–71.
- 52 Schnebel B, Gwin JT, Anderson S, et al. In vivo study of head impacts in football: a comparison of national collegiate athletic association division I versus high school impacts. *Neurosurgery* 2007;60:490–6.
- 53 Brolinson PG, Manoogian S, McNeely D, et al. Analysis of linear head accelerations from collegiate football impacts. *Curr Sports Med Rep* 2006;5:23–8.
- 54 McAllister TW, Ford JC, Ji S, et al. Maximum principal strain and strain rate associated with concussion diagnosis correlates with changes in corpus callosum white matter indices. *Ann Biomed Eng* 2012;40:127–40.
- 55 Funk JR, Rowson S, Daniel RW, et al. Validation of concussion risk curves for collegiate football players derived from HITS data. *Ann Biomed Eng* 2012;40:79–89.
- 56 Duhaime AC, Beckwith JG, Maerlender AC, et al. Spectrum of acute clinical characteristics of diagnosed concussions in college Athletes wearing instrumented helmets: clinical article. *J Neurosurg* 2012;117:1092–9.
- 57 Duma SM, Rowson S. Every Newton hertz: a macro to micro approach to investigating brain injury. *Conf Proc IEEE Eng Med Biol Soc* 2009;2009:1123–6.
- 58 King D, Hume PA, Brughelli M, et al. Instrumented mouthguard acceleration analyses for head impacts in amateur rugby union players over a season of matches. *Am J Sports Med* 2015;43:614–24.
- 59 Hernandez F, Wu LC, Yip MC, et al. Six Degree-of-Freedom measurements of human mild traumatic brain injury. *Ann Biomed Eng* 2015;43:1918–34.
- 60 Wilcox BJ, Beckwith JG, Greenwald RM, et al. Biomechanics of head impacts associated with diagnosed concussion in female collegiate ice hockey players. *J Biomech* 2015;48:2201–4.
- 61 Beckwith JG, Chu JJ, Crisco JJ, et al. Severity of head impacts resulting in mild traumatic brain injury. *Am Soc Biomech* 2009:1144.



What is the definition of sports-related concussion: a systematic review

Paul McCrory, Nina Feddermann-Demont, Jiri Dvorák, J David Cassidy, Andrew McIntosh, Pieter E Vos, Ruben J Echemendia, Willem Meeuwisse and Alexander A Tarnutzer

Br J Sports Med 2017 51: 877-887
doi: 10.1136/bjsports-2016-097393

Updated information and services can be found at:
<http://bjsm.bmj.com/content/51/11/877>

	<i>These include:</i>
References	This article cites 55 articles, 7 of which you can access for free at: http://bjsm.bmj.com/content/51/11/877#BIBL
Email alerting service	Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>